For Research Use Only

PCSK9 Recombinant antibody

Catalog Number:84172-4-RR




Basic Information	Catalog Number: 84172-4-RR	GenBank Accession Number: NM_174936.4	Purification Method: Protein A purification					
	Concentration:	GenelD (NCBI):	CloneNo.:					
	1000 µg/ml	255738	241395G1					
	Source:	UNIPROT ID:	Recommended Dilutions:					
	Rabbit Isotype: IgG	Q8NBP7 Full Name: proprotein convertase subtilisin/kexin type 9	WB 1:500-1:2000 IF/ICC 1:200-1:800					
				Calculated MW: 74 kDa Observed MW: 72-78 kDa, 62 kDa				
						Applications	Tested Applications:	Positive
		WB, IF/ICC, FC (Intra), ELISA	WB : HepG2 cells,					
		Species Specificity: human	IF/ICC : HepG2 cells,					
		Background Information	Proprotein convertase subtilisin/kexin type 9 (PCSK9) is a crucial protein governing the circulating levels of low density lipoprotein-cholesterol (LDL-C), by virtue of its pivotal role in the degradation of the LDL receptor (LDLR). PCSK9 is expressed in the kidney and lung. It is synthesized as a 72 kDa immature precursor that undergoes autocatalytic cleavage in the endoplasmic reticulum to generate a 63 kDa mature protein. The cleaved N-terminal fragment remains associated with the mature protein and is necessary for its secretion, allowing it to circulate in the blood. The ability of PCSK9 to regulate a diverse group of cell-surface proteins hinted that it might also be able to influence additional membrane proteins that are important in anti-tumour immune responses. Targeting PCSK9 to treat cancer is also attractive because two neutralizing antibodies against it, evolocumab and alirocumab, have already been approved for human clinical use to lower cholesterol levels. (PMID: 30522786, PMID: 22493497)					
Storage	Storage: Store at -20°C. Stable for one yea Storage Buffer: PBS with 0.02% sodium azide an Aliquoting is unnecessary for -20	d 50% glycerol						

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data






HepG2 cells were subjected to SDS PAGE followed by western blot with 84172-4-RR (PCSK9 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.

Immunofluorescent analysis of (4% PFA) fixed HepC2 cells using PCSK9 antibody (84172-4-RR, Clone: 241395G1) at dilution of 1:400 and CoraLite@488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



1x10^6 HeLa cells were intracellularly stained with 0.25 ug PCSK9 Recombinant antibody (8&172-4-R, Clone:241395G1) and CoraLite®488-Conjugated Goat Anti-Rabbit 1gG(H+L) (SA00013-2)(red), or 0.25 ug Rabbit 1gG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLI) kinetic assays of 84172-4-RR against Human PCSK9 were performed. The affinity constant is 0.80 nM.