

For Research Use Only

Ki-67 Recombinant antibody

Catalog Number: 84192-4-RR **2 Publications**



Basic Information

Catalog Number: 84192-4-RR	GenBank Accession Number: NM_002417	Purification Method: Protein A purification
Concentration: 1000 ug/ml	GeneID (NCBI): 4288	CloneNo.: 241499E7
Source: Rabbit	UNIPROT ID: P46013	Recommended Dilutions: WB 1:5000-1:50000 IHC 1:1000-1:4000 IF-P 1:50-1:500 IF/ICC 1:200-1:800
Isotype: IgG	Full Name: antigen identified by monoclonal antibody Ki-67	
	Calculated MW: 359 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, IF-P, ELISA	Positive Controls:
Cited Applications: IHC, IF	WB : HeLa cells, HepG2 cells, HEK-293 cells, MCF-7 cells, Ramos cells
Species Specificity: human	IHC : human tonsillitis tissue, human skin cancer tissue, Human Spleen tissue, Human renal cancer, human malignant melanoma tissue, human prostate cancer tissue, human Breast cancer tissue
Cited Species: human	IF-P : human lung cancer tissue, human thyroid cancer tissue
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	IF/ICC : HeLa cells, MCF-7 cells, U2OS cells, HepG2 cells, A431 cells, A549 cells

Background Information

The Ki-67 protein (also known as MKI67) is a cellular marker for proliferation. Ki67 is present during all active phases of the cell cycle (G1, S, G2 and M), but is absent in resting cells (G0). Cellular content of Ki-67 protein markedly increases during cell progression through S phase of the cell cycle. Therefore, the nuclear expression of Ki67 can be evaluated to assess tumor proliferation by immunohistochemistry. It has been demonstrated to be of prognostic value in breast cancer. In head and neck cancer, several studies have reported an association between high proliferative activity and poorer prognosis.

Notable Publications

Author	Pubmed ID	Journal	Application
Ming Zhou	39831459	Cell Biol Int	IHC
Penghui Ye	39787763	Int Immunopharmacol	IF


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 84192-4-RR (Ki-67 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 84192-4-RR (Ki-67 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human lung cancer tissue using Ki-67 antibody (84192-4-RR, Clone: 241499E7) at dilution of 1:200 and Multi-rAb CoraLite® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), ICAM-1/CD54 antibody (60299-1-Ig, Clone: 2F9A8, red). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Ki-67 antibody (84192-4-RR, Clone: 241499E7) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human lung cancer tissue using Ki-67 antibody (84192-4-RR, Clone: 241499E7) at dilution of 1:200 and Multi-rAb CoraLite® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 84192-4-RR (Ki-67 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.