## For Research Use Only

## Geminin Recombinant antibody

Catalog Number:84199-2-RR



**Purification Method:** 

Protein A purfication

Recommended Dilutions:

CloneNo.:

240835B5

**Basic Information** 

Catalog Number: GenBank Accession Number: 84199-2-RR BC005185

 Concentration:
 GeneID (NCBI):

 1000 μ g/ml
 51053

 Source:
 UNIPROT ID:

 Rabbit
 075496
 WB 1:500-1:2000

 Isotype:
 Full Name:
 IHC 1:250-1:1000

 IgG
 geminin, DNA replication inhibitor
 IF/ICC 1:50-1:500

Immunogen Catalog Number:Calculated MW:AG107624 kDa

Observed MW: 28-29 kDa

**Applications** 

Tested Applications: WB, IHC, IF/ICC, ELISA Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

human

Positive Controls:

WB: Jurkat cells, HEK-293 cells
IHC: human colon cancer tissue,
IF/ICC: HeLa cells, HepG2 cells

## **Background Information**

GMNN, also designated geminin, contains 212 amino acids and has a destruction box sequence (RRTLKVIQP). GMNN participates in inhibiting DNA replication by preventing the incorporation of MCM complex into the pre-replication complex (pre-RC). It is degraded during the metaphase-anaphase transition of cell cycle's mitotic phase, which permits replication in the succeeding cell cycle. GMNN has a broad sedimentation profile ranging from about 25 kDa to 90 kDa, with a major peak at 30 kDa. Scanning of the signals shows that discrete peaks corresponding to the apparent mass of 42.5 kDa and 66 kDa are present. The dimer of geminin(50 kDa) can also be detected. (PMID: 15313623)

Storage


Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data




Jurkat cells were subjected to SDS PAGE followed by western blot with 84199-2-RR (GMNN antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



HEK-293 cells were subjected to SDS PAGE followed by western blot with 84199-2-RR (GMNN antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 84199-2-RR (Geminin antibody) at dilution of 1:500 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Geminin antibody (84199-2-RR, Clone: 240835B5) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Geminin antibody (84199-2-RR, Clone: 240835B5) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



Biolayer interferometry (BLI) kinetic assays of 84199-2-RR against Human Geminin were performed. The affinity constant is below 1 pM.