For Research Use Only

MLH1 Recombinant antibody, PBS Only (Capture)

Catalog Number:84208-2-PBS



Basic Information

Catalog Number:

GenBank Accession Number: BC006850 Purification Method:
Protein A purification

241472D1

84208-2-PBS Size:

GeneID (NCBI):

CloneNo.:

1 mg/ml Source: 4292 UNIPROT ID:

UNIPROT ID: P40692

Rabbit Isotype:

Full Name: mutL homolog 1, colon cancer,

Immunogen Catalog Number:

nonpolyposis type 2 (E. coli)

AG2319 Calculated MW:

756 aa, 85 kDa Observed MW: 85-100 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, Cytometric bead array, Indirect

ELISA

Species Specificity:

human

Background Information

MLH1, also named as COCA2, belongs to the DNA mismatch repair mutL/hexB family. It heterodimerizes with PMS2 to form MutL alpha which is a component of the post-replicative DNA mismatch repair system (MMR). MutL alpha (MLH1-PMS2) interacts physically with the clamp loader subunits of DNA polymerase III, suggesting that it may play a role to recruit the DNA polymerase III to the site of the MMR. MLH1 also implicated in DNA damage signaling, a process which induces cell cycle arrest and can lead to apoptosis in case of major DNA damages. MLH1 heterodimerizes with MLH3 to form MutL gamma which plays a role in meiosis.(PMID: 16873062, PMID: 18206974) Defects in MLH1 are the cause of hereditary non-polyposis colorectal cancer type 2 (HNPCC2). Defects in MLH1 are a cause of mismatch repair cancer syndrome (MMRCS). Defects in MLH1 are a cause of Muir-Torre syndrome (MTS). Defects in MLH1 are a cause of susceptibility to endometrial cancer. Western blot analysis with an MLH1 antibody detected a 85-100 kDa band.

Storage

Storage:


Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Ruffor.


Storage Buffer:

PBS Only

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 84208-2-RR (MLH1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84208-2-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 84208-2-RR (MLH1 antibody) at dilution of 1:250 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 84208-2-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed HeLa cells using MLH1 antibody (84208-2-RR, Clone: 241472D1) at dilution of 1:300 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 84208-2-PBS in a different storage buffer formulation.




IP result of anti-MLH1 (IP:84208-2-RR, 4ug; Detection:84208-2-RR 1:1000) with Hela cells lysate 1200 ug. This data was developed using the same antibody clone with 84208-2-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed HeLa cells using MLH1 antibody (84208-2-RR, Clone: 241472D1) at dilution of 1:300 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red). This data was developed using the same antibody clone with 84208-2-PBS in a different storage buffer formulation.



Cytometric bead array standard curve of MP01140-1, MLH1 Recombinant Matched Antibody Pair, PBS Only. Capture antibody: 84208-2-PBS. Detection antibody: 84208-1-PBS. Standard: Ag2319. Range: 0.313-40 ng/mL



Biolayer interferometry (BLI) kinetic assay of 84208-2-PBS against Human MLH1 was performed. The affinity constant is below 1 pM.



Biolayer interferometry (BLI) kinetic assays of 84208-2-RR against Human MLH1 were performed. The affinity constant is below 1 pM.