For Research Use Only


VDAC2 Recombinant antibody, PBS Only (Capture) Catalog Number:84225-2-PBS



Catalog Number: 84225-2-PBS	GenBank Accession Number: BC000165	Purification Method: Protein A purification
Size: 1 mg/ml	GenelD (NCBI): 7417	CloneNo.: 241569G11
Source: Rabbit	UNIPROT ID: P45880	
Isotype: IgG	Full Name: voltage-dependent anion channel 2	
Immunogen Catalog Number: AG2266	Calculated MW: 294 aa, 32 kDa	
	Observed MW: 31-33 kDa	
Tested Applications: WB, FC (Intra), Cytometric bead array, Indirect ELISA		
Species Specificity: human, mouse, rat		
VDACs (Voltage Dependent Anion selective Channels), also known as mitochondrial porins, are a family of pore- forming proteins discovered in the mitochondrial outer membrane. Mammals show a conserved genetic organization of the VDAC genes. It's reported that the amount of VDAC transcripts in liver is usually lower than in the other tissues. VDAC2 and especially VDAC3 are highly expressed in testis, while mouse VDAC1 is poorly expressed in this tissue. (PMID: 22020053)		
Storage: Store at -80°C. The product is shipped with ice pa Storage Buffer: PBS Only	icks. Upon receipt, store it immediatel	ly at −80°C
	84225-2-PBS Size: 1 mg/ml Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG2266 Tested Applications: WB, FC (Intra), Cytometric bead an Species Specificity: human, mouse, rat VDACs (Voltage Dependent Anior forming proteins discovered in the organization of the VDAC genes. I the other tissues. VDAC2 and espe expressed in this tissue. (PMID: 22 Storage: Storage Storage Buffer:	84225-2-PBS BC000165 Size: GeneID (NCBI): 1 mg/ml 7417 Source: UNIPROT ID: Rabbit P45880 Isotype: Full Name: IgG voltage-dependent anion channel Immunogen Catalog Number: Calculated MW: AG2266 294 aa, 32 kDa Observed MW: 31-33 kDa Tested Applications: WB, FC (Intra), Cytometric bead array, Indirect ELISA Species Specificity: human, mouse, rat VDACs (Voltage Dependent Anion selective Channels), also known as a forming proteins discovered in the mitochondrial outer membrane. Mar organization of the VDAC genes. It's reported that the amount of VDAC the other tissues. VDAC2 and especially VDAC3 are highly expressed in expressed in this tissue. (PMID: 22020053) Storage: Storage Liss hipped with ice packs. Upon receipt, store it immediated Storage Buffer:


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




100 - 80 Relative cell Number 60 40 20


Various lysates were subjected to SDS PAGE followed by western blot with 84225-2-RR (VDAC2 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84225-2-PBS in a different storage buffer formulation.



1x10^6 HepG2 cells were intracellularly stained with 0.25 ug Vdac2 Recombinant Antibody (84225-2-RR, Clone:241569G11) and APC-Conjugated Goat Anti-Rabbit $\lg G(H+L)(red)$, or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 84225-2-PBS in a different storage buffer formulation.



Cytometric bead array standard curve of MP01162-1, VDAC2 Recombinant Matched Antibody Pair, PBS Only. Capture antibody: 84225-2-PBS. Detection antibody: 84225-1-PBS. Standard: Ag2266. Range: 1.563-200 ng/mL



Biolayer interferometry (BLI) kinetic assays of 84225-2-RR against Human VDAC2 were performed. The affinity constant is 0.48 nM.