

For Research Use Only

CD98/SLC3A2 Recombinant antibody, PBS Only

Catalog Number: 84387-6-PBS



Basic Information

Catalog Number:

84387-6-PBS

Size:

1 mg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_002394.5

GeneID (NCBI):

6520

UNIPROT ID:

P08195-1

Full Name:

solute carrier family 3 (activators of
dibasic and neutral amino acid
transport), member 2

Calculated MW:

68kDa

Observed MW:

68-125 kDa

Purification Method:

Protein A purification

CloneNo.:

241755A11

Applications

Tested Applications:

WB, IF/ICC, Indirect ELISA

Species Specificity:

human

Background Information

CD98 is a 125-kDa type II membrane glycoprotein heterodimer composed of a 40-kDa non-glycosylated light chain (amino-acid transporter) and an 85-kDa glycosylated heavy chain (CD98hc) (PMID: 11507094). CD98hc also interacts with certain integrin β -subunits to regulate cell migration, survival, proliferation, and adhesion/polarity. CD98hc is overexpressed on the cell surface of many cancers and increased CD98hc expression is associated with the development and progression of tumors (PMID: 25084765). Molecular orientation studies have demonstrated that the C-terminal tail of human CD98 (hCD98), which contains a PDZ-binding domain, is extracellular (PMID: 18625289).

Storage

Storage:

Store at -80°C .

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84387-6-RR (CD98 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84387-6-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CD98 antibody (84387-6-RR, Clone: 241755A11) at dilution of 1:250 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). This data was developed using the same antibody clone with 84387-6-PBS in a different storage buffer formulation.