For Research Use Only

ADAMTS2 Recombinant antibody

Isotype:

Catalog Number:84476-6-RR



Purification Method:

Protein A purfication

Recommended Dilutions:

WB 1:5000-1:50000 IF/ICC 1:125-1:500

CloneNo.:

241860B8

Basic Information

Catalog Number: GenBank Accession Number:

84476-6-RR NM_014244 GeneID (NCBI): Size: 1000 µg/ml 9509 **UNIPROT ID:** Source: Rabbit 095450

ADAM metallopeptidase with thrombospondin type 1 motif, 2 Immunogen Catalog Number:

AG34957 Calculated MW:

135 kDa Observed MW: 110~130 kDa

Full Name:

Applications

Tested Applications: Positive Controls:

WB, IF/ICC, ELISA WB: A431 cells, COLO 205 cells, PC-3 cells, HT-1080 Species Specificity:

cells

IF/ICC: A431 cells,

Background Information

A disintegrin and metalloproteinase with thrombospondin motifs 2 (ADAMTS-2) is a metalloproteinase that plays a key role in the processing of fibrillar procollagen precursors into mature collagen molecules by excising the aminopropeptide. ADAMTS-2 can, like the procollagen C-proteinases, be regulated by transforming growth factor- β 1 (TGF- β 1), with implications for mechanisms whereby this growth factor effects net increases in formation of $extracellular\,matrix.\,Mutations\,in\,the\,ADAMTS-2\,gene\,cause\,dermatosparaxis\,in\,human\,(dermatosparactic\,type\,of\,allow)$ Ehlers-Danlos syndrome, also known as Ehlers-Danlos syndrome type VIIC). Human ADAMTS-2 has eight potential sites for Asn-linked glycosylation and other mammals (PMID: 16046392, 12646579).

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84476-6-RR (ADAMTS2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed A431 cells using ADAMT52 antibody (84476-6-RR, Clone: 241860B8) at dilution of 1:250 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).



Biolayer interferometry (BLI) kinetic assays of 84476-6-RR against Human ADAMTS2 were performed. The affinity constant is 0.472 nM.