For Research Use Only

CD3 Recombinant antibody

Catalog Number:84505-4-RR



Purification Method:

Protein A purfication

Recommended Dilutions:

WB 1:5000-1:50000 IHC 1:500-1:2000

IF/ICC 1:200-1:800

CloneNo.:

242012G2

Basic Information

Catalog Number: 84505-4-RR

Size: 1000 µg/ml

Source: Rabbit Isotype: IgG

Full Name: CD3e molecule, epsilon (CD3-TCR

GenBank Accession Number:

BC049847

916

GeneID (NCBI):

ENSEMBL Gene ID:

ENSG00000198851

UNIPROT ID:

P07766

complex)

Calculated MW: 207 aa, 23 kDa Observed MW: 23 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: Jurkat cells, MOLT-4 cells IHC: human tonsillitis tissue,

IF/ICC : Jurkat cells,

Background Information

CD3 is a complex of proteins that directly associates with the T cell receptor (TCR). The TCR/CD3 complex of T-lymphocytes consists of either a TCR alpha/beta or TCR gamma/delta heterodimer coexpressed at the cell surface with the invariant subunits of CD3 labeled gamma, delta, epsilon, zeta, and eta. The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction. CD3 is considered to be a pan-T cell marker for detection of normal and neoplastic T cells. This antibody is directed against the epsilon chain of human CD3 molecule.

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 84505-4-RR (CD3E antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 84505-4-RR (CD3 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 84505-4-RR (CD3 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed Jurkat cells using CD3 antibody (84505-4-RR, Clone: 242012G2) at dilution of 1:400 and CoraLite@488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).