For Research Use Only

## Acetyl-Histone H2B (Lys20) Recombinant antibody, PBS Only

Catalog Number:84549-1-PBS



**Basic Information** 

Catalog Number: 84549-1-PBS

Concentration: 1 mg/ml

Source: Rabbit Isotype:

G histone cluster 2, H2be

Calculated MW: 14 kDa Observed MW: 15 kDa

GenBank Accession Number:

BC005827

8349

GeneID (NCBI):

**UNIPROT ID:** 

Q16778
Full Name:

Purification Method: Protein A purfication

CloneNo.: 241204H8

**Applications** 

Tested Applications:

WB, IF/ICC, Dot Blot, Indirect ELISA

Species Specificity: human, mouse

## **Background Information**

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS only, pH7.3


## Selected Validation Data




Dot blot analysis was used to confirm the specificity of 84549-1-RR Acetyl-Histone H2B (Lys20) antibody. Peptides were spotted onto NC and probed with antibody at 1 µg/ml. The amount of peptide (ug/ml.) spotted is indicated next to each row. This data was developed using the same antibody clone with 84549-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed Trichostatin A treated NIH/3T3 cells using Acetyl-Histone H2B (Lys20) antibody (84549-1-RR, Clone: 241204H8) at dilution of 1:1000 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (5A00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 84549-1-PBS in a different storage buffer formulation.



Trichostatin A treated and untreated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 84549-1-RR (Acetyl-Histone H2B (Lys20) antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Alpha Tubulin (66031-1-Ig) antibody as a loading control. This data was developed using the same antibody clone with 84549-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed Trichostatin A treated HeLa cells using Acetyl-Histone H2B (Lys20) antibody (84549-1-RR, Clone: 241204H8) at dilution of 1:1000 and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 84549-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Acetyl-Histone H2B (Lys20) antibody (84549-1-RR, Clone: 241204H8) at dilution of 1:400 and Coralite®488-Conjugated Goat Anti-Rabbit 1gG(H+L) (SA00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 84549-1-PBS in a different storage buffer formulation.