

For Research Use Only

Acetyl-Histone H2B (Lys20) Recombinant antibody

Catalog Number: 84549-1-RR



Basic Information

Catalog Number:

84549-1-RR

Concentration:

1000 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC005827

GeneID (NCBI):

8349

UNIPROT ID:

Q16778

Full Name:

histone cluster 2, H2be

Calculated MW:

14 kDa

Observed MW:

15 kDa

Purification Method:

Protein A purification

CloneNo.:

241204H8

Recommended Dilutions:

WB 1:500-1:2000

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IF/ICC, Dot Blot, ELISA

Species Specificity:

human, mouse

Positive Controls:

WB : Trichostatin A treated NIH/3T3 cells,

IF/ICC : HeLa cells, Trichostatin A treated HeLa cells,
Trichostatin A treated NIH/3T3 cells

Background Information

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Dot blot analysis was used to confirm the specificity of 84549-1-RR Acetyl-Histone H2B (Lys20) antibody. Peptides were spotted onto NC and probed with antibody at 1 µg/ml. The amount of peptide (µg/mL) spotted is indicated next to each row.




Trichostatin A treated and untreated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 84549-1-RR (Acetyl-Histone H2B (Lys20) antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Alpha Tubulin (66031-1-IG) antibody as a loading control.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Acetyl-Histone H2B (Lys20) antibody (84549-1-RR, Clone: 241204H8) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed Trichostatin A treated NIH/3T3 cells using Acetyl-Histone H2B (Lys20) antibody (84549-1-RR, Clone: 241204H8) at dilution of 1:1000 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed Trichostatin A treated HeLa cells using Acetyl-Histone H2B (Lys20) antibody (84549-1-RR, Clone: 241204H8) at dilution of 1:1000 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).