For Research Use Only

MGEA5 Recombinant antibody

Catalog Number:84722-4-RR



Purification Method:

Protein A purfication

Recommended Dilutions:

WB 1:5000-1:50000 IHC 1:125-1:500

IF/ICC 1:200-1:800

CloneNo.:

242263A6

Basic Information

Catalog Number: 84722-4-RR

GeneID (NCBI): 10724

Source: 060502 Full Name:

Calculated MW: 103 kDa

Observed MW: 130-140 kDa

Size: 1000 µg/ml

Rabbit Isotype:

Immunogen Catalog Number: AG6405

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA

Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

GenBank Accession Number:

BC039583

UNIPROT ID:

meningioma expressed antigen 5

(hyaluronidase)

Positive Controls:

WB: HeLa cells, JAR cells

IHC: human rectal cancer tissue,

IF/ICC: HepG2 cells,

Background Information

Human meningioma-expressed antigen 5 (MGEA5) has two putative domains including protein O-GlcNAcase domain and histone acetyltransferase domain, therefore it is often called bifunctional protein NCOAT. Three isoforms of MEGE5 are produced by alternative splicing. MGEA5 was found to be regulated to reduce the state of glycosylation of transcitpional activators while increasing the acetylation of histones to allow for the concerted activation of eukaryotic gene transcription, for instance, acetylation of Lys8 of histone H4 and Lys 14 of histone H3 $are \ resulted \ from \ acetyl transferase \ activity. \ In \ addition, single \ nucleotide \ polymorphism \ in \ MGE5A \ is \ associated$ with type 2 diabetes in Mexican Americans. Two bands at 130kDa and 75 kDa could be detected using the present rabbit polyclonal antibody 14711-1-AP, which is consistent with results in a related reference (PubMed:11341771).


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 84722-4-RR (MGEA5 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human rectal cancer tissue slide using 84722-4-RR (MGEA5 antibody) at dilution of 1:250 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using MGEA5 antibody (84722-4-RR, Clone: 242263A6) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



Biolayer interferometry (BLI) kinetic assays of 84722-4-RR against Human MGEA5 were performed. The affinity constant is below 1 pM.