For Research Use Only

APOE Recombinant antibody

Catalog Number:84775-6-RR





Basic Information	Catalog Number: 84775-6-RR	GenBank Accession Number: BC003557	Purification Method: Protein A purfication	
	Size: 1000 ug/ml	GenelD (NCBI): 348	CloneNo.: 242310B3	
	Source: Rabbit	UNIPROT ID: P02649	Recommended Dilutions: WB 1:2000-1:10000	
	Isotype: IgG	Full Name: Apolipoprotein E		
		Observed MW: 34-38 kDa		
Applications	Tested Applications: WB, FC (Intra), ELISA Species Specificity: human	Positive Controls:		
		WB : Hept	WB : HepG2 cells, HuH-7 cells, human placenta	
Background Information	The apolipoprotein E (APOE) is a 299-amino acid polypeptide that mediates the binding, internalization, and catabolism of lipoprotein particles, and also serves as a ligand for the LDL (apo B/E) receptor and for the specific apo-E receptor (chylomicron remnant) of hepatic tissues. The very strong association of the APOE 4 allele with AD risk and its role in the accumulation of amyloid β in brains of people and animal models solidify the biological relevance of APOE isoforms but do not provide mechanistic insight (PMID: 9831633).			
Storage	Storage: Store at -20°C. Stable for one Storage Buffer: PBS with 0.02% sodium azid			
	A11 6 6	Aliquoting is unnecessary for -20°C storage		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1x10^6 HepG2 cells were intracellularly stained with 0.25 ug APOE Recombinant antibody (84775-6-RR, Clone:242310B3) and PE-Conjugated Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLI) kinetic assays of 84775-6-RR against Human APOE were performed. The affinity constant is below 1 pM.

Various lysates were subjected to SDS PAGE followed by western blot with 84775-6-RR (APOE antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.