For Research Use Only

GOLIM4 Recombinant antibody, PBS Only




Catalog Number:84818-5-PBS

Basic Information	Catalog Number: 84818-5-PBS	GenBank Accession Number: NM_014498	Purification Method: Protein A purfication
	Size:	GenelD (NCBI):	CloneNo.:
	1 mg/ml	27333	242323F4
	Source: Rabbit	UNIPROT ID: 000461	
	Isotype:Full Name:IgGgolgi integral membrane protein 4		4
	Immunogen Catalog Number: AG34962	Calculated MW: 82 kDa	
	Observed MW: 130-140 kDa		
Applications	Tested Applications: WB, IF/ICC, FC (Intra), Indirect ELI	SA	
	Species Specificity: human, mouse		
Background Informatior	Golgi integral membrane protein 4 (GOLIM4) is a 130-kDa membrane protein that is integrated into the cis/medial Golgi apparatus, which is a kind II Golgi protein. GOLIM4 is a Golgi protein that circulates between Golgi apparatus and endosomes, and its shuttle is pH sensitive(PMID:30068697).		
Storage	Storage: Store at -80°C.	cks. Upon receipt, store it immediatel	vat-80°C
	Storage Buffer: PBS Only	eks. opometerpt, store it inimediater	y at-60 C


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84818-5-RR (GOLIM4 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84818-5-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed U2OS cells using GOLIM4 antibody (84818-5-RR, Clone: 242323F4) at dilution of 1:500 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 84818-5-PBS in a different storage buffer formulation.

1.5

(mu

Response

 $K_D = 9.26 \times 10^{-10}$ $K_{off} = 1.61 \times 10^{-4}$ $K_{on} = 1.74 \times 10^{5}$ 200 nM


100 nM 50 nM

25 nM


12.5 nM

6.25 nM

3.125 nM



1x10^{^6} U2OS cells were intracellularly stained with 0.25 ug GOLIM4 Recombinant antibody (84818-5-RR, Clone:242323F4) and APC-Conjugated Goat Anti-Rabbit IgG(H+L) (red), or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 84818-5-PBS in a different storage buffer



1x10^6 HeLa cells were intracellularly stained with 0.25 ug GOLIM4, Recombinant antibody (84818-5-RR, Clone:242323F4) and APC-Conjugated Goat Anti-Rabbit IgG(H+L) (red), or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 84818-5-PBS in a different storage buffer 0.0 100 200 300 400 Time(s) Biolayer interferometry (BLL) kinetic assays of

84818-5-RR against Human GOLIM4 were performed. The affinity constant is 0.926 nM.