

For Research Use Only

# Mouse P62/SQSTM1 Recombinant antibody, PBS Only (Capture/Detector)

Catalog Number: 84826-1-PBS



## Basic Information

Catalog Number:

84826-1-PBS

Concentration:

1 mg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG35064

GenBank Accession Number:

NM\_011018

GeneID (NCBI):

18412

UNIPROT ID:

Q64337

Full Name:

sequestosome 1

Calculated MW:

48 kDa

Observed MW:

62 kDa

Purification Method:

Protein A purification

CloneNo.:

241992C4

## Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), IP, Cytometric bead array, Sandwich ELISA, Indirect ELISA, Sample test

Species Specificity:

human, mouse, rat

## Background Information

Sequestosome 1 (SQSTM1/p62) is a multifunctional adaptor protein implicated in selective autophagy, cell signaling pathways, and tumorigenesis. p62 has been implicated in shuttling ubiquitinated and aggregated proteins for autophagic degradation. p62 is degraded during the autophagic process, which makes its intracellular level a marker for autophagy progression. p62 is at the cross-roads of several signaling pathways including Ras/ Raf/ MAPK and NF- $\kappa$ B and plays important role in cancer.

## Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 84826-1-RR (P62/SQSTM1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84826-1-PBS in a different storage buffer formulation.




NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 84826-1-RR (P62/SQSTM1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84826-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 84826-1-RR (P62/SQSTM1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84826-1-PBS in a different storage buffer formulation.




Sandwich ELISA standard curve of MP01608-2, Mouse P62,Sqstm1 Recombinant Matched Antibody Pair - PBS only. 84826-1-PBS was coated to a plate as the capture antibody and incubated with serial dilutions of standard Ag35064. 84826-2-PBS was HRP conjugated as the detection antibody. Range: 0.781-50 ng/mL




The mean P62,Sqstm1 concentration was determined to be 257.13 ng/mL in mouse brain tissue extract based on a 1.6 mg/mL extract load and 598.93 ng/mL in NIH/3T3 cell extract based on a 1.5 mg/mL extract load.




IP result of anti-P62/SQSTM1 (IP:84826-1-RR, 4ug; Detection:84826-1-RR 1:4000) with NIH/3T3 cells lysate 1920 ug. This data was developed using the same antibody clone with 84826-1-PBS in a different storage buffer formulation.






Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 84826-1-RR (P62/SQSTM1 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 84826-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 84826-1-RR (P62/SQSTM1 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 84826-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using P62/SQSTM1 antibody (84826-1-RR, Clone: 241992C4) at dilution of 1:500 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 84826-1-PBS in a different storage buffer formulation.



1x10<sup>6</sup> NIH/3T3 cells were intracellularly stained with 0.25 ug P62/SQSTM1 Recombinant antibody (84826-1-RR, Clone:241992C4) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone



Cytometric bead array standard curve of MP01608-1, MOUSE P62,Sqstm1 Recombinant Matched Antibody Pair, PBS Only. Capture antibody: 84826-2-PBS. Detection antibody: 84826-1-PBS. Standard: Ag35064. Range: 0.625-80 ng/mL

Biolayer interferometry (BLI) kinetic assays of 84826-1-RR against Mouse P62,SQSTM1 were performed. The affinity constant is 0.203 nM.