For Research Use Only

Phospho-POLR2A (Ser5) Recombinant antibody

Catalog Number:84834-1-RR



Basic Information

Catalog Number: 84834-1-RR

Rabbit P24928 Isotype: Full Name:

gG polymerase (RNA) II (DNA directed)

polypeptide A, 220kDa Calculated MW:

GenBank Accession Number:

1970 aa, 217 kDa Observed MW: 270 kDa

BC137231

Applications

Tested Applications: WB, FC (Intra), ELISA Species Specificity: Positive Controls:

WB: HeLa cells, λ phosphatase treated HeLa cells

Purification Method:

Protein A purfication

WB 1:5000-1:50000

Recommended Dilutions:

CloneNo.:

241914E5

Background Information

RNA Pol II consists of 12 subunits and is responsible for the expression of mRNA and lincRNA genes within cells. The subunits of RNA polymerases have been shown to play specific roles in Pol II CTD dephosphorylation, tissue-specific transcription, mRNA decay, and translation. Pol II transcription involves initiation through recruitment and CTD serine 5 phosphorylation, elongation with CTD serine 2 phosphorylation, and termination. (PMID: 38842922)

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Non-treated HeLa cells, phosphatase inhibitor treated HeLa cells and $^\lambda$ phosphatase treated HeLa cells were subjected to SDS PAGE followed by western blot with 84834-1-RR (Phospho-POLR2A (Ser5) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Lamin B1 (66095-1-lg) antibody as a loading control.



1X10^6 HeLa cells untreated (dashed lines) or treated with $^{\lambda}$ phosphatase which intracellularly stained with 0.13 ug Phospho-POLR2A (Ser5) Recombinant antibody (84834-1-RR, Clone:241914E5) and Coralite®488-Conjugated Goat Anti-Rabbit 1gG(H+L) (5A00013-2)(red), or 0.13 ug Rabbit 1gG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.