For Research Use Only

CD3 Gamma Recombinant antibody

Catalog Number:84836-2-RR



Purification Method:

Protein A purfication

Recommended Dilutions:

WB 1:5000-1:50000 IHC 1:500-1:2000

CloneNo.:

241849D4

Basic Information

Catalog Number: 84836-2-RR

Concentration: Genel D (NCBI): 1000 ug/ml 917

Source: UNIPROT ID:
Rabbit P09693
Isotype: Full Name:

gG CD3g molecule, gamma (CD3-TCR

Immunogen Catalog Number: complex)

EG2147 Calculated MW: 182 aa, 20 kDa

Observed MW: 22 kDa

BC113830

GenBank Accession Number:

Applications

Tested Applications: WB, IHC, ELISA

Species Specificity: human Note-IHC: suggested antigen retrieval with

TE buffer pH 9.0; (*) Alternatively, antigen

retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: Jurkat cells, MOLT-4 cells IHC: human tonsillitis tissue,

Background Information

CD3 is a complex of proteins that directly associates with the T cell receptor (TCR). The TCR/CD3 complex of T-lymphocytes consists of either a TCR alpha/beta or TCR gamma/delta heterodimer coexpressed at the cell surface with the invariant subunits of CD3 labeled gamma, delta, epsilon, zeta, and eta. The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction. CD3 is considered to be a pan-T cell marker for detection of normal and neoplastic T cells.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various cells were subjected to SDS PAGE followed by western blot with 84836-2-RR (CD3 Gamma antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 84836-2-RR (CD3 Gamma antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 84836-2-RR (CD3 Gamma antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).