## For Research Use Only

## **HGD** Recombinant antibody

Catalog Number:84931-5-RR



**Purification Method:** 

Protein A purfication

WB 1:5000-1:50000

Recommended Dilutions:

CloneNo.:

242485E11

**Basic Information** 

Catalog Number: 84931-5-RR

Concentration:

BC020792 GeneID (NCBI): 3081 **UNIPROT ID:** 

1000  $\,\mu$  g/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG9544

**Tested Applications:** 

WB, ELISA Species Specificity: human, rat

GenBank Accession Number:

Q93099 Full Name:

Calculated MW: 37 kDa, 50 kDa Observed MW: 50 kDa

homogentisate 1,2-dioxygenase (homogentisate oxidase)

Positive Controls:

WB: A549 cells, HepG2 cells

## **Background Information**

Homogentisate1,2-dioxygenase (HGD), also named as HGO, is a mononuclear Fe(II)-dependent oxygenase that catalyzes the third step in the pathway for the catabolism of tyrosine, the conversion of homogentisate (HG) to the conversion of homogeneous catalyzes the third step in the pathway for the catabolism of tyrosine, the conversion of homogeneous catalyzes the third step in the pathway for the catabolism of tyrosine, the conversion of homogeneous catalyzes the third step in the pathway for the catabolism of tyrosine, the conversion of homogeneous catalyzes the catabolism of thmaleylacetoacetate (MAA) and it can exsit as a dimer or trimer(PMID:14678794). HGD consists of a single type of subunit with no intermolecular disulfide bridges and requires Fe2+ as a cofactor(PMID: 7705358). Defects in HGD are the cause of alkaptonuria (AKU)(PMID:10594001).

Storage

**Applications** 


Store at -20°C. Stable for one year after shipment.


Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data







Biolayer interferometry (BLL) kinetic assays of 84931-5-RR against Human HGD were performed. The affinity constant is 1.38 nM.