

For Research Use Only

HNRNPU Recombinant antibody

Catalog Number: 84964-4-RR



Basic Information

Catalog Number:

84964-4-RR

Concentration:

1000 μ g/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG6141

GenBank Accession Number:

BC003621

GeneID (NCBI):

3192

UNIPROT ID:

Q00839

Full Name:

heterogeneous nuclear
ribonucleoprotein U (scaffold
attachment factor A)

Calculated MW:

825 aa, 91 kDa

Observed MW:

120 kDa

Purification Method:

Protein A purification

CloneNo.:

242609C9

Recommended Dilutions:

WB 1:5000-1:50000

Applications

Tested Applications:

WB, FC (Intra), ELISA

Species Specificity:

human, mouse

Positive Controls:

WB : HEK-293 cells, Jurkat cells, HeLa cells, HepG2
cells, MCF-7 cells, NIH/3T3 cells,

Background Information

HNRNPU, also named as HNRPU, SAFA, and U21.1, is a 825 amino acid protein, which is extensively phosphorylated. HNRNPU as a component of ribonucleosomes localized in cytoplasmic mRNP granules containing untranslated mRNAs. HNRNPU is a component of the CRD-mediated complex that promotes MYC mRNA stabilization. HNRNPU Plays a role in the circadian regulation of the core clock component ARNTL/BMAL1 transcription. The calculated molecular weight of HNRNPU is 90 kDa, but the 120 kDa HNRNPU protein is an abundant nucleoplasmic phosphoprotein that can be crosslinked to pre-mRNA in intact cells by UV light (PMID: 1628625).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84964-4-RR (HNRNPU antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



1×10^6 HepG2 cells were intracellularly stained with 0.25 μ g HNRNPU Recombinant antibody (84964-4-RR, Clone:242609C9) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2) (red), or 0.25 μ g Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).