For Research Use Only

MEK1/2 Recombinant antibody, PBS Only (Capture/Detector)

Catalog Number:84967-1-PBS



Purification Method:

Protein A purification

CloneNo.:

242614G3

Basic Information

Catalog Number: 84967-1-PBS Concentration:

1 mg/ml
Source:
Rabbit
Isotype:

gG mitogen-activated protein kinase

Immunogen Catalog Number: kinase 2

AG1481 Calculated MW:

44 kDa Observed MW: 44 kDa

BC018645

5605

P36507 Full Name:

GeneID (NCBI):

UNIPROT ID:

GenBank Accession Number:

Applications

Tested Applications:

WB, IF/ICC, FC (Intra), Cytometric bead array, Sandwich ELISA, Indirect ELISA, Sample test

Species Specificity: human, mouse, rat

Background Information

MEK2(MAPK/ERK kinase 2) is also named as MAP2K2, MKK2, PRKMK2 and belongs to the MAP kinase kinase subfamily. MAPKK is itself dependent on Ser/Thr phosphorylation for activity catalyzed by MAP kinase kinase kinases (RAF or MEKK1). MEK1 and MEK2 are closely related, dual-specificity tyrosine/threonine protein kinases found in the Ras/Raf/MEK/ERK mitogen-activated protein kinase (MAPK) signaling pathway. 84967-1-RR can recognize both the MEK2 and MEK1.

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS Only


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 84967-1-RR (MEK1/2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84967-1-PBS in a different storage buffer formulation.




The mean MEK2 concentration was determined to be 45.61 ng/mL in A549 cell extract based on a 1.50 mg/mL extract load and 44.07 ng/mL in HeLa cell extract based on a 1.00 mg/mL extract load.




1x10^6 Jurkat cells were intracellularly stained with 0.25 ug MEK1/2 Recombinant Antibody (84967-1-RR, Clone:242614G3) and Coralite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 84967-1-PBS in a different storage buffer




Immunofluorescent analysis of (4% PFA) fixed Jurkat cells using MEK1/2 antibody (84967-1-RR, Clone: 242614G3) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). This data was developed using the same antibody clone with 84967-1-PBS in a different storage buffer formulation.




Sandwich ELISA standard curve of MP01692-1, Human MEK2 Recombinant Matched Antibody Pair -PBS only. 84967-2-PBS was coated to a plate as the capture antibody and incubated with serial dilutions of standard Ag1481. 84967-1-PBS was HRP conjugated as the detection antibody. Range: 0.781-50 ng/mL.



Cytometric bead array standard curve of MP01692-1, MEK2 Recombinant Matched Antibody Pair, PBS Only. Capture antibody: 84967-2-PBS. Detection antibody: 84967-1-PBS. Standard: Ag1481. Range: 0.625-80 ng/mL



Cytometric bead array standard curve of MP01692-2, MEK2 Recombinant Matched Antibody Pair, PBS Only. Capture antibody: 84967-1-PBS. Detection antibody: 84967-3-PBS. Standard: Ag1481. Range: 0.625-80 ng/mL



Biolayer interferometry (BLI) kinetic assays of 84967-1-RR against Human MEK1/2 were performed. The affinity constant is below 1 pM.