For Research Use Only

ALR Recombinant antibody

Catalog Number:84972-4-RR




| Basic Information | Catalog Number: 84972-4-RR | GenBank Accession Number: BC028348 | Purification Method: Protein A purification |
|------------------------|---|---|--|
| | Concentration: 1000 µg/ml | GenelD (NCBI): 2671 | CloneNo.: 242557G3 |
| | Source: Rabbit | UNIPROT ID: P55789 | Recommended Dilutions: WB 1:500-1:2000 |
| | lsotype: lgG Immunogen Catalog Number: AG1840 | Full Name: growth factor, augmenter of liver regeneration Calculated MW: 15 kDa, 23 kDa | |
| | | | |
| | | Applications | Tested Applications: WB, FC (Intra), ELISA Species Specificity: human, mouse, rat |
| Background Information | GFER (FAD-linked sulfhydryl oxidase) is also named as ALR, HERV1, HPO. It plays an important role in the disulfide relay system (DRS) in human mitochondria. The GFER gene codes for 2 distinct isoforms that are probably synthesized from the same mRNA with the use of different initiation codons. The long isoform (205 amino acids, 23/21 kD) is located mainly in the mitochondrial intermembrane space and exists under nonreducing and nondenaturing conditions as a homodimer and a heterodimer. The shorter isoform (125 amino acids, 15 kD), which lacks 80 amino acids at its N terminus compared to the longer isoform, is present predominantly in the nucleus (PMID: 19409522, 24880092, 21152698). | | |
| Storage | Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20° | 50% glycerol pH 7.3. | |


For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84972-4-RR (GFER antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



1x10^6 HepG2 cells were intracellularly stained with 0.25 ug ALR Recombinant antibody (84972-4-RR, Clone:242557G3) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2) (red), or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLL) kinetic assays of 84972-4-RR against Human ALR were performed. The affinity constant is 0.23 nM.