For Research Use Only

SNRNP200 Recombinant antibody, PBS Only




Catalog Number:84992-5-PBS

| Basic Information | Catalog Number: 84992-5-PBS | GenBank Accession Number: BC001417 | Purification Method: Protein A purification |
|------------------------|---|--|--|
| | Concentration: 1 mg/ml | GeneID (NCBI): 23020 | CloneNo.: 242550D10 |
| | Source: Rabbit | UNIPROT ID: O75643 | |
| | Isotype: IgG Immunogen Catalog Number: AG20748 | Full Name: small nuclear ribonucleoprotein 200kDa (U5) | |
| | | Calculated MW: 494 aa, 57 kDa | |
| | | Observed MW: 200-245 kDa | |
| Applications | Tested Applications: WB, Indirect ELISA | | |
| | Species Specificity: human | | |
| Background Information | SNRNP200 (small nuclear ribonucleoprotein 200kDa (U5)), also known as HELIC2, ASCC3L1 or BRR2, is a 2,136 amino acid protein that localizes to the nucleus and contains two SEC63 domains, two helicase C-terminal domains and two helicase ATP-binding domains. Existing as multiple alternatively spliced isoforms, HELIC2 functions as an RNA helicase that is thought to promote specific RNA-RNA conformational changes which are important in the second step of RNA splicing. The gene encoding HELIC2 maps to human chromosome 2, which houses over 1,400 genes and comprises nearly 8% of the human genome. Harlequin icthyosis, a rare and morbid skin deformity, is associated with mutations in the chromosome 2-localized ABCA12 gene, while the lipid metabolic disorder sitosterolemia is associated with defects in the ABCG5 and ABCG8 genes, hich also map to chromosome 2. The observed molecular weight of SNRNP200 is 200-245kd. | | |
| Storage | Storage: Store at -80°C. The product is shipped with ice pa Storage Buffer: PBS only, pH7.3 | cks. Upon receipt, store it immediatel | yat-80°C |

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84992-5-RR (SNRNP200 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84992-5-PBS in a different storage buffer formulation.