

For Research Use Only

CD155/PVR Recombinant antibody

Catalog Number: 85006-1-RR



Basic Information

Catalog Number:

85006-1-RR

Size:

1000 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_027514.2

GeneID (NCBI):

52118

UNIPROT ID:

Q8K094

Full Name:

poliovirus receptor

Calculated MW:

45 kDa

Observed MW:

70-80 kDa

Purification Method:

Protein A purification

CloneNo.:

242243G9

Recommended Dilutions:

WB 1:5000-1:50000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

mouse

Positive Controls:

WB : RAW 264.7 cells, mouse pancreas tissue, mouse small intestine tissue, mouse placenta tissue

Background Information

CD155, also known as the poliovirus receptor (PVR) or Nectin-5, is one of the nectin-like family members, belonging to the Ca²⁺ independent immunoglobulin superfamily (IgSF) in cell adhesion molecules (PMID: 37660884). CD155 is a type I transmembrane protein composed of three extracellular immunoglobulin-like domains, a transmembrane domain, and a cytoplasmic tail. It is involved in cell-to-cell and cell-to-ECM adhesion (PMID: 32019260; 11437656). CD155 acts as a ligand for immune receptors, including TIGIT, CD96 and CD226, which are expressed on T cells and NK cells. CD155 interacts with these receptors to modulate immune responses, such as T cell activation and NK cell-mediated cytotoxicity (PMID: 37660884). CD155 serves as the entry receptor for poliovirus and thereby mediates susceptibility to poliovirus infection (PMID: 15034010).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 85006-1-RR (CD155/PVR antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.