

## ACSS1 Recombinant antibody, PBS Only

Catalog Number: 85102-4-PBS

## Basic Information

## Catalog Number:

85102-4-PBS

## Concentration:

1 mg/ml

## Source:

Rabbit

## Isotype:

IgG

## Immunogen Catalog Number:

AG10853

## GenBank Accession Number:

BC039261

## GeneID (NCBI):

84532

## UNIPROT ID:

Q9NUB1

## Full Name:

acyl-CoA synthetase short-chain  
family member 1

## Calculated MW:

689 aa, 75 kDa

## Observed MW:

75 kDa

## Purification Method:

Protein A purification

## CloneNo.:

242361C6

## Applications

## Tested Applications:

WB, FC (Intra), Indirect ELISA

## Species Specificity:

human, mouse, rat

## Background Information

The ACSS (acetyl-CoA synthetase) enzyme is the sole known mammalian enzyme that can catalyze the conversion of free acetate into acetyl coenzyme A (acetyl-CoA). The three known isoforms of human ACSS are termed ACSS1, ACSS2, and ACSS3. The main substrate of ACSS1 and ACSS2 is acetate, while the preferential substrate of ACSS3 is propionate. Two acetate related enzymes, ACSS1 (GeneID: 84532) and ACSS2 (GeneID: 55902) differ in their tissue distribution and subcellular localization. On the one hand, as a mitochondrial matrix enzyme, ACSS1 is expressed mainly in cardiac and skeletal muscle as well as brown adipose tissue. On the other hand, as a nuclear and cytoplasmic enzyme, ACSS2 is strongly expressed in the liver, kidney and heart and moderately expressed in the brain and testis. ACSS2 participates in lipid synthesis and facilitates protein acetylation by generating acetyl-CoA, while ACSS1 is involved in acetate oxidation. The functional differences in these enzymes involve energy production through the tricarboxylic acid (TCA) cycle. Due to its more thorough utilization of intracellular acetate, ACSS2 is expressed in almost all cell types under different physiological conditions.

## Storage

## Storage:

Store at -80°C.

**The product is shipped with ice packs. Upon receipt, store it immediately at -80°C**

## Storage Buffer:

PBS only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 85102-4-RR (ACSS1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 85102-4-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 85102-4-RR (ACSS1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 85102-4-PBS in a different storage buffer formulation.



1x10<sup>6</sup> HeLa cells were intracellularly stained with 0.25 ug ACS1 Recombinant antibody (85102-4-RR, Clone:242361C6) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 85102-4-PBS in a different storage buffer