For Research Use Only

MCT2/SLC16A7 Recombinant antibody, PBS Only

Catalog Number:85145-3-PBS



Purification Method:

Protein A purification

CloneNo.:

242811F6

Basic Information

Catalog Number: 85145-3-PBS

Concentration:
1 mg/ml
Source:
Rabbit

Isotype: IgG

Immunogen Catalog Number:

AG14201

Applications

Tested Applications: WB, Indirect ELISA Species Specificity:

Background Information

SLC16A7, also named MCT2, is a proton-coupled monocarboxylate transporter catalyzing the rapid transport across the plasma membrane of many monocarboxylates. It has been reported that SLC16A7 is the primary monocarboxylate transporter in Purkinje cells and can be detected in the brain, cerebellum, liver, and kidney, but not in vascular smooth muscle cells, pericytes, endothelium, heart, lung, or stomach. There are some isoforms of SLC16A7 with bands around 52 kDa and 40-45 kDa in SDS-PAGE. The 80 kDa protein might be a dimer, similar to the papers published. (PMID: 9482213, 15749979, 11291733, 16901736)

GenBank Accession Number:

solute carrier family 16, member 7 (monocarboxylic acid transporter 2)

BC030693

9194

O60669 Full Name:

GeneID (NCBI):

UNIPROT ID:

Calculated MW: 478 aa, 52 kDa Observed MW: 40-52 kDa

Storage

T: 4006900926

Storage: Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer:

PBS only, pH7.3


For technical support and original validation data for this product please contact:

E: Proteintech-CN@ptglab.com


W: ptgcn.cor

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 85145-3-RR (SLC16A7 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 85145-3-PBS in a different storage buffer formulation.



Biolayer interferometry (BLL) kinetic assays of 85145-3-RR against Human MCT2/SLC 16A7 were performed. The affinity constant is 37.0 nM.