

For Research Use Only

# DNAJC7 Recombinant antibody

Catalog Number: 85157-4-RR



## Basic Information

<b>Catalog Number:</b> 85157-4-RR	<b>GenBank Accession Number:</b> BC011837	<b>Purification Method:</b> Protein A purification
<b>Concentration:</b> 1000 µg/ml	<b>GeneID (NCBI):</b> 7266	<b>CloneNo.:</b> 242875B3
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> Q99615	<b>Recommended Dilutions:</b> WB 1:5000-1:50000
<b>Isotype:</b> IgG	<b>Full Name:</b> DnaJ (Hsp40) homolog, subfamily C, member 7	IHC 1:50-1:500
<b>Immunogen Catalog Number:</b> AG1572	<b>Calculated MW:</b> 56 kDa	IF/ICC 1:200-1:800
	<b>Observed MW:</b> 56 kDa	

## Applications

<b>Tested Applications:</b> WB, IHC, IF/ICC, FC (Intra), ELISA	<b>Positive Controls:</b> WB : HeLa cells, HepG2 cells, A549 cells, HEK-293 cells, Jurkat cells, mouse testis tissue, rat testis tissue
<b>Species Specificity:</b> human, mouse, rat	<b>IHC :</b> mouse brain tissue,
<b>Note-IHC:</b> <i>suggested antigen retrieval with <b>TE buffer pH 9.0</b>; (*) Alternatively, antigen retrieval may be performed with <b>citrate buffer pH 6.0</b></i>	<b>IF/ICC :</b> HepG2 cells,

## Background Information

DnaJC7, otherwise referred to as Tpr2 or CCRP, is a conserved, type of J protein that is highly expressed within the brain. DnaJC7 has been found to modulate the flow of substrates from Hsp70 to Hsp90, providing a control mechanism in proteins such as glucocorticoids and progesterone receptor.

## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol  
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 85157-4-RR (DNAJC7 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using DNAJC7 antibody (85157-4-RR, Clone: 242875B3 ) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



1x10<sup>6</sup> HeLa cells were intracellularly stained with 0.25 ug DNAJC7 Recombinant antibody (85157-4-RR, Clone:242875B3) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2) (red), or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLI) kinetic assays of 85157-4-RR against Human DNAJC7 were performed. The affinity constant is 0.767 nM.