

For Research Use Only

# ICOSLG/B7-H2/CD275 Recombinant monoclonal antibody, PBS Only

Catalog Number:85173-5-PBS



## Basic Information

<b>Catalog Number:</b> 85173-5-PBS	<b>GenBank Accession Number:</b> BC064637	<b>Purification Method:</b> Protein A purification
<b>Source:</b> Rabbit	<b>GeneID (NCBI):</b> 23308	<b>CloneNo.:</b> 242241G9
<b>Isotype:</b> IgG	<b>UNIPROT ID:</b> O75144	
<b>Immunogen Catalog Number:</b> EG0008	<b>Full Name:</b> inducible T-cell co-stimulator ligand	
	<b>Calculated MW:</b> 33 kDa, 34 kDa	
	<b>Observed MW:</b> 65 kDa	

## Applications

**Tested Applications:**  
WB, Indirect ELISA

**Species Specificity:**  
human

## Background Information

ICOSLG, also known as B7H2 or GL50, is a type I transmembrane glycoprotein belonging to the B7 ligand family. ICOSLG is extensively expressed on professional antigen-presenting cells including B cells, macrophages, and dendritic cells, as well as non-lymphoid cells including mesenchymal stem cells, endothelial cells, fibroblasts, and tumor cells (PMID: 33756276; 35800767). It is a specific ligand for the T-cell-specific cell surface receptor ICOS and acts as a co-stimulatory molecule (PMID: 11023515; 11007762). The interaction of ICOSLG and ICOS plays important roles in the activation, proliferation, differentiation, and cytokine production of T cells as well as in the antibody secretion from B cells during secondary immune responses (PMID: 21851236).

## Storage

**Storage:**  
Store at -80°C.  
**The product is shipped with ice packs. Upon receipt, store it immediately at -80°C**

**Storage Buffer:**  
PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)

W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data



JAR cells were subjected to SDS PAGE followed by western blot with 85173-5-RR (ICOSLG antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 85173-5-PBS in a different storage buffer formulation.