For Research Use Only

TRPC4 Recombinant antibody

Catalog Number:85334-2-RR



Purification Method:

Protein A purification

Recommended Dilutions:

WB 1:2000-1:10000

CloneNo.:

242444F8

Basic Information

Catalog Number: GenBank Accession Number: 85334-2-RR BC104725

 $\begin{tabular}{llll} Concentration: & GenelD (NCBI): \\ 1000 ~~\mu ~g/ml & 7223 \\ Source: & UNIPROT ID: \\ \end{tabular}$

Source: UNIPROT ID:
Rabbit Q9UBN4
Isotype: Full Name:

IgG transient receptor potential cation channel, subfamily C, member 4

AG15698 Calculated MW: 977 aa, 112 kDa
Observed MW:

Observed MW: 140 kDa

Applications

Tested Applications: WB, ELISA

Species Specificity:

human

Positive Controls:

WB: A549 cells, HEK-293 cells, SH-SY5Y cells

Background Information

TRPC4 (Transient Receptor Potential Paradigm 4) is a non-selective calcium channel protein widely found in the nervous system and cardiovascular system. TRPC4 regulates intracellular calcium levels through the activation of signaling pathways mediated by Gq/11 and Gi/o-coupled receptors, and is involved in neurotransmission, neuronal excitability, and vascular endothelial cell function. In the nervous system, TRPC4 is associated with neuropsychiatric disorders such as depression and anxiety, and its inhibitors show antidepressant and anxiolytic potential. In the cardiovascular system, TRPC4 affects vascular permeability and diastolic function by regulating calcium ion inward flow and plays an important role in hypoxia-induced pulmonary hypertension. In addition, TRPC4 is involved in thermoregulation, sensing changes in internal temperature to regulate the drop in body temperature. aberrant expression of TRPC4 is associated with a variety of diseases, making it a potential therapeutic target!

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 85334-2-RR (TRPC4 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.