For Research Use Only

VAMP8 Recombinant antibody

Catalog Number:85400-4-RR



Basic Information

Catalog Number: GenBank Accession Number: 85400-4-RR BC001634

Concentration: Genel D (NCBI): CloneNo.: 1000 μ g/ml 8673 242830B11

Source: UNIPROT ID: Recommended Dilutions:
Rabbit Q9BV40 WB 1:5000-1:50000
Isotype: Full Name: IHC 1:1000-1:4000
IgG vesicle-associated membrane protein IF/ICC 1:200-1:800

Immunogen Catalog Number: 8 (endobrevin)

AG7903 Calculated MW:

11 kDa
Observed MW:
15 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA

Species Specificity: human, mouse, rat

buffer pH 6.0

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate Positive Controls:

WB: THP-1 cells, HL-60 cells, A431 cells, HeLa cells, rat

Purification Method:

Protein A purification

kidney tissue, mouse kidney tissue

IHC: human placenta tissue,

IF/ICC : THP-1 cells,

Background Information

VAMP8, also named as endobrevin, is a member of the vesicle-associated membrane protein (VAMP)/synaptobrevin family and the SNARE (soluble NSF-attachment protein receptor) superfamily. Characterized by a common sequence called the SNARE motif, SNARE proteins are involved in membrane fusion and vesicular transport (PMID: 11252968). VAMP8 is involved in autophagy through the direct control of autophagosome membrane fusion with the lysososome membrane. It is required for dense-granule secretion in platelets and plays a role in regulated enzyme secretion in pancreatic acinar cells. VAMP8 is also involved in the abscission of the midbody during cell division, which leads to completely separate daughter cells.

Storage


Storage:

Storage Ruffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 85400-4-RR (VAMP8 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 85400-4-RR (VAMP8 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 85400-4-RR (VAMP8 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed THP-1 cells using VAMP8 antibody (85400-4-RR, Clone: 242830B11) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).



Biolayer interferometry (BLL) kinetic assays of 85400-4-RR against Human VAMP8 were performed. The affinity constant is 0.631 nM.