

For Research Use Only

p21 Recombinant monoclonal antibody

Catalog Number: 85435-2-RR



Basic Information

Catalog Number: 85435-2-RR	GenBank Accession Number: NM_001111099	Purification Method: Protein A purification
Source: Rabbit	GeneID (NCBI): 12575	CloneNo.: 242478H2
Isotype: IgG	UNIPROT ID: P39689	Recommended Dilutions: WB: 1:500-1:2000 IF/ICC: 1:500-1:2000
Immunogen Catalog Number: AG28249	Full Name: cyclin-dependent kinase inhibitor 1A (P21)	
	Calculated MW: 18 kDa	
	Observed MW: 15-18 kDa	

Applications

Tested Applications: WB, IF/ICC, ELISA	Positive Controls:
Species Specificity: mouse	WB : NIH/3T3 cells, Neuro-2a cells, DC2.4 cells, RAW 264.7 cells
	IF/ICC : NIH/3T3 cells,

Background Information

CDKN1A (p21, CIP1, WAF1) is a cyclin-dependent kinase inhibitor. CDKN1A binds to and inhibits the activity of cyclin-CDK2 or -CDK4 complexes, and thus functions as a regulator of cell cycle progression at the G1 phase. The expression of CDKN1A is induced by wild-type but not mutant p53 protein, through which CDKN1A mediates the p53-dependent cell cycle G1 phase arrest in response to a variety of stress stimuli. CDKN1A can interact with proliferating cell nuclear antigen (PCNA), and plays a regulatory role in S phase DNA replication and DNA damage repair. CDKN1A was reported to be specifically cleaved by CASP3-like caspases, which thus leads to a dramatic activation of CDK2, and may be instrumental in the execution of apoptosis following caspase activation.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.

Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 85435-2-RR (Cdkn1a antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using Cdkn1a antibody (85435-2-RR, Clone: 242478H2) at dilution of 1:1000 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



Biolayer interferometry (BLI) kinetic assays of 85435-2-RR against Mouse p21 were performed. The affinity constant is below 1 pM.