For Research Use Only

## NOV/CCN3 Recombinant antibody

Catalog Number:85731-1-RR



Basic Information	Catalog Number: 85731-1-RR	GenBank Accession Number: NM_010930.4	Purification Method: Protein A purification
	Concentration: 1000 µg/ml	Genel D (NCBI): 18133	CloneNo.: 250054B2
	Source: Rabbit	UNIPROT ID: Q64299	Recommended Dilutions: WB 1:1000-1:4000
	Isotype: IgG	Full Name: nephroblastoma overexpressed gene	
	Immunogen Catalog Number: EG2856	Calculated MW: 39 kDa	
		Observed MW: 46 kDa	
Applications	Tested Applications: WB, ELISA	WB : A172 cells, HEK-293T cells, mouse brain tissu ecificity: HeLa cells, mouse thymus tissue, Caco-2 cells, rat	
	Species Specificity: human, mouse, rat		
	CCN family member 3 (CCN3), also known as Nephroblastoma overexpressed (NOV) protein, is a member of the CCN protein family (PMID: 1309586). CCN3 is predominantly expressed in the cytoplasm, nucleus, and extracellular space. As a subset of extracellular matrix proteins, it can directly bind to the integrin receptor family on the cell surface and participate in signal transduction between the cell surface and the extracellular matrix structure (PMID: 37378812). CCN3 exhibits anti-proliferative activity in normal cells and most tumor cells (PMID: 17340618). CCN3 co-localizes with connexin 43 (C×43) in the plasma membrane, and these two proteins physically interact to inhibit cell growth (PMID: 15213231). CCN3 can induce myofibroblast differentiation through the Notch pathway (PMID: 12050162). CCN can promote adhesion of endothelial cells, vascular smooth muscle cells, and fibroblasts through heparan sulfate proteoglycans (HSPG) (PMID: 30158025).		
Background Information	CCN protein family (PMID: 130958 space. As a subset of extracellular surface and participate in signal t 37378812). CCN3 exhibits anti-pr co-localizes with connexin 43 (C× cell growth (PMID: 15213231). CC 12050162). CCN can promote adh	86). CCN3 is predominantly expressed matrix proteins, it can directly bind to ransduction between the cell surface a oliferative activity in normal cells and 43) in the plasma membrane, and the N3 can induce myofibroblast different esion of endothelial cells, vascular sm	I in the cytoplasm, nucleus, and extracellular o the integrin receptor family on the cell and the extracellular matrix structure (PMID: d most tumor cells (PMID: 17340618). CCN3 se two proteins physically interact to inhibit iation through the Notch pathway (PMID:

 For technical support and original validation data for this product please contact:

 T: 4006900926
 E: Proteintech-CN@ptglab.com
 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 85731-1-RR (Nov antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 85731-1-RR (Nov antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Mouse brain tissue was subjected to SDS PAGE followed by western blot with 85731-1-RR (Nov antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.