For Research Use Only

NMDAR1/GRIN1 Recombinant antibody

Catalog Number:85973-3-RR



Basic Information

Catalog Number: 85973-3-RR

 Concentration:
 GeneID (NCBI):

 1000 μ g/ml
 2902

 Source:
 UNIPROT ID:

Rabbit Q05586

Isotype: Full Name:
IgG glutamate receptor, ionotropic, N-

Immunogen Catalog Number: methyl D-aspartate 1

AG26093 Calculated MW: 105 kDa

Observed MW: 116-120 kDa

NM 000832

GenBank Accession Number:

Purification Method:

Protein A purification CloneNo.:

Recommended Dilutions:

WB: 1:1000-1:4000 IP: 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate

250535B1

Applications

Tested Applications: WB, IP, ELISA

Species Specificity: human, mouse, rat

Positive Controls:

WB: mouse brain tissue, rat brain tissue

IP: mouse brain tissue,

Background Information

GRIN1 encodes subunit 1 of the N-methyl-D-aspartate (NMDA) receptor, which is a heteromeric glutamate-gated calcium ion channel essential for synaptic function in the brain (PMID: 25864721, PMID: 25864721). NMDARs play important roles in normal brain development and function, such as synaptic plasticity, neural development, learning and memory (PMID: 20716669). NMDAR dysfunction has been associated with several neurological disorders including Parkinson, Alzheimer and Huntington diseases. Disrupted motor learning and long-term synaptic plasticity in mice lacking NMDAR1 in the striatum (PMID: 17015831).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 85973-3-RR (NMDAR1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



IP result of anti-NMDAR1/GRIN1 (IP:85973-3-RR, 4ug; Detection:85973-3-RR 1:1000) with mouse brain tissue lysate 1720 ug.



Biolayer interferometry (BLI) kinetic assays of 85973-3-RR against Human NMDAR1/GRIN1 were performed. The affinity constant is below 1 pM.