

CD8a Recombinant antibody

Catalog Number: 85977-3-RR

Basic Information

Catalog Number:

85977-3-RR

Concentration:

1000 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

EG3265

GenBank Accession Number:

NM_001081110.2

GeneID (NCBI):

12525

UNIPROT ID:

P01731

Full Name:

CD8 antigen, alpha chain

Calculated MW:

27 kDa

Observed MW:

32-35 kDa

Purification Method:

Protein A purification

CloneNo.:

242879G4

Recommended Dilutions:

WB: 1:1000-1:6000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

mouse

Positive Controls:

WB : mouse thymus tissue,

Background Information

CD8 is a transmembrane glycoprotein that is predominantly expressed on the surface of cytotoxic T cells, and can also be found on natural killer cells, cortical thymocytes, and dendritic cells. CD8 is composed of two disulfide-linked chains and can be present as a homodimer of CD8 α or as a heterodimer of CD8 α and CD8 β (PMID: 3264320; 8253791). The majority of class I-restricted T cells express mostly the CD8 α β heterodimer while CD8 α α homodimers alone have been found on some gut intraepithelial T cells, on some T cell receptor (TCR) γ δ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4⁻/CD8⁺ T cells (PMID: 1968084).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Mouse thymus tissue was subjected to SDS PAGE followed by western blot with 85977-3-RR (CD8A antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.