## For Research Use Only

## BRD2 Recombinant antibody

Catalog Number:85990-2-RR



**Basic Information** 

Catalog Number: GenBank Accession Number: 85990-2-RR BC063840

gG bromodomain containing 2

Immunogen Catalog Number:Calculated MW:AG17589836 aa, 92 kDa

Observed MW: 100-110 kDa

Full Name:

Purification Method:

Protein A purification

CloneNo.: 250453G5

Recommended Dilutions: WB: 1:5000-1:50000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IF/ICC: 1:250-1:1000

Applications

Tested Applications: WB, IF/ICC, IP, ELISA Species Specificity:

human, mouse, rat

Isotype:

Positive Controls:

WB: HeLa cells, Jurkat cells, A549 cells, K-562 cells,

C2C12 cells, PC-12 cells

IP: HeLa cells,

IF/ICC: Jurkat cells,

## **Background Information**

BRD2 (Bromodomain-containing protein 2), also known as RING3, is a member of the Bromodomain and Extra-Terminal (BET) family of proteins. These proteins contain two bromodomains and one extra-terminal domain. BRD2 primarily localizes to the nucleus and functions as an atypical protein kinase. Its main role involves the activation of E2F1 and E2F2 proteins, facilitating the production of proteins essential for the G1/S phase transition of the cell cycle. BRD2 specifically recognizes acetylated histones through its bromodomains, particularly acetylated lysine residues on histones H3 and H4. This interaction allows BRD2 to recruit various transcription factors, co-activators, and repressors, thereby modulating transcriptional activity. BRD2 plays a crucial role in chromatin remodeling by recruiting proteins such as histone deacetylases. It is highly expressed during neural tube development and closure, making it essential for neurogenesis and embryogenesis.(PMID: 33462181, PMID: 37926722)

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 85990-2-RR (BRD2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



IP result of anti-BRD2 (IP:85990-2-RR, 4ug; Detection:85990-2-RR 1:3000) with HeLa cells lysate 1600 ug.



Immunofluorescent analysis of (4% PFA) fixed Jurkat cells using BRD2 antibody (85990-2-RR, Clone: 250453G5) at dilution of 1:500 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).