

For Research Use Only

CA8 Recombinant antibody

Catalog Number: 86004-2-RR



Basic Information

Catalog Number:	86004-2-RR	GenBank Accession Number:	BC015531	Purification Method:	Protein A purification
Concentration:	1000 µg/ml	GeneID (NCBI):	767	Clone No.:	250429C4
Source:	Rabbit	UNIPROT ID:	P35219	Recommended Dilutions:	WB: 1:5000-1:50000
Isotype:	IgG	Full Name:	carbonic anhydrase VIII		
Immunogen Catalog Number:	AG3068	Calculated MW:	290 aa, 32 kDa		
		Observed MW:	32-35 kDa		

Applications

Tested Applications:	WB, ELISA	Positive Controls:	WB: HEK-293 cells, mouse brain tissue, Jurkat cells, K-562 cells, rat brain tissue
Species Specificity:	human, mouse, rat		

Background Information

The CA8 (CARP) gene encodes carbonic anhydrase VIII, which is part of a family of zinc metalloenzyme. CA8 has a central carbonic anhydrase motif, but it lacks carbonic anhydrase activity due to absence of catalytic zinc coordinating residues (PMID:2121526). CARP is a novel IP3R1-binding protein, and is expressed in Purkinje cells abundantly. CA8 is co-localized with IP3R1 in Purkinje cells and it binds to IP3R1, reducing the affinity of the receptor for its ligand, IP3 (PMID:12611586). Defects in CA8 are the cause of cerebellar ataxia mental retardation and dysequilibrium syndrome type 3 (CMARQ3) (PMID:19461874).

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 86004-2-RR (CA8 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 86004-2-RR (CA8 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLI) kinetic assays of 86004-2-RR against Human CA8 were performed. The affinity constant is below 1 pM.