For Research Use Only

GIRK2 Recombinant antibody, PBS Only

Catalog Number:86016-1-PBS


Antibodies | ELISA kits | Proteins Uni-rAb www.ptglab.com


Basic Information	Catalog Number: 86016-1-PBS	GenBank Accession Number: BC101547	Purification Method: Protein A purification	
	Concentration: 1 mg/ml	GenelD (NCBI): 3763	CloneNo.: 250605D1	
	Source: Rabbit	UNIPROT ID: P48051		
	lsotype: IgG Immunogen Catalog Number: AG16344	Full Name: potassium inwardly-rectifying channel, subfamily J, member 6		
		Calculated MW: 423 aa, 48 kDa		
		Observed MW: 45-48 kDa		
Applications	Tested Applications: WB, Indirect ELISA			
	Species Specificity: human, mouse, rat			
Background Information	GIRK2 (also known as Kir3.2), encoded by KCNJ6 gene, is a member of the G protein-coupled inwardly-rectifying potassium channel (GIRK, Kir3) family of inward rectifier potassium channels. GIRK channels are activated following stimulation of G protein-coupled receptors (PMID: 26422984). They play important roles in regulating cellular excitabilities in the heart and brain (PMID: 31043612). Mutations in KCNJ6 gene are associated with Keppen-Lubinsky Syndrome (PMID: 25620207).			
Storage	Storage: Store at -80°C. The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer: PBS only, pH7.3			

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 86016-1-RR (KCNJ6 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 86016-1-PBS in a different storage buffer formulation.



Biolayer interferometry (BLI) kinetic assays of 86016-1-RR against Human GIRK2 were performed. The affinity constant is 0.372 nM.

Various lysates were subjected to SDS PAGE followed by western blot with 86016-1-RR (KCNJ6 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 86016-1-PBS in a different storage buffer formulation.