For Research Use Only

AIF Recombinant antibody

Catalog Number:86156-3-RR



Basic Information	Catalog Number: 86156-3-RR	GenBank Accession Number: BC 111065	Purification Method: Protein A purification					
	Concentration: 1000 µg/ml	GenelD (NCBI): 9131	CloneNo.: 250772F8					
	Source: Rabbit	UNIPROT ID: O95831	Recommended Dilutions: WB: 1:5000-1:50000					
	lsotype: IgG Immunogen Catalog Number: AG13268	Full Name: apoptosis-inducing factor, mitochondrion-associated, 1						
		Calculated MW: 609 aa, 66 kDa						
		Observed MW: 67 kDa						
Applications	Tested Applications:	Positive Controls: WB : HeLa cells, HEK-293T cells, Jurkat cells, K-562 cells, NIH/3T3 cells, C2C12 cells, C6 cells						
	WB, ELISA Species Specificity: human, mouse, rat							
Background Information	Apoptosis-inducing factor (AIF) is one of the mitochondrial proteins to be released into the cytosol during apoptosis, and it is discovered as the first protein that regulates caspase-independent apoptosis(PMID:20494118). AIF is encoded as a 67 kDa protein that contains a mitochondrial localization signal (MLS) in the N-terminus. It is cleaved from the 62 kDa to the 57 kDa form following ischemic injury and translocated from the mitochondria to the nucleus in a calpain-dependent manner(PMID:19332058).							
Storage	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol, pH7.3 Aliquoting is unnecessary for -20°C storage							

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data

	HeLa	HEK-25	13T Iurkal	K.552	NIH/3T	3 02012	c6	
180 kDa→								
140 kDa \rightarrow								
$100 \text{ kDa} \rightarrow$								
75 kDa→								
60 kDa→	-	-	-	-	-		-	← AIF
G								
$45 \ kDa {\rightarrow}$								
$35 \ \mathrm{kDa}{ ightarrow}$								
ŏ								
25 kDa→								

Various lysates were subjected to SDS PAGE followed by western blot with 86156-3-RR (AIF antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.