

For Research Use Only

BRD7 Recombinant antibody

Catalog Number: 86176-2-RR



Basic Information

Catalog Number:

86176-2-RR

Concentration:

1000 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG1461

GenBank Accession Number:

BC050728

GeneID (NCBI):

29117

UNIPROT ID:

Q9NPI1

Full Name:

bromodomain containing 7

Calculated MW:

651 aa, 74 kDa

Observed MW:

74-78 kDa

Purification Method:

Protein A purification

CloneNo.:

250776B2

Recommended Dilutions:

WB: 1:2000-1:10000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

human, mouse

Positive Controls:

WB : Jurkat cells, mouse lung tissue, PC-3 cells, SH-SY5Y cells, COLO 320 cells

Background Information

BRD7 (bromodomain containing 7), is a subunit of poly-bromo-associated BRG1-associated factor (PBAF)-specific Swi/Snf chromatin remodeling complexes. BRD7 is proposed as a tumor suppressor protein, since its identification as an important component of both functional p53 and BRCA1 (breast cancer 1, early onset) pathways. BRD7 could down-regulate the GSK3β phosphotransferase activity as dephosphorylation of GSK3β at 'Tyr-216', thus activate the Wnt signaling pathway in a DVL1-dependent manner.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 86176-2-RR (BRD7 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



PC-3 cells were subjected to SDS PAGE followed by western blot with 86176-2-RR (BRD7 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



mouse lung tissue were subjected to SDS PAGE followed by western blot with 86176-2-RR (BRD7 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.