

For Research Use Only

# TORC1/CRTC1 Recombinant antibody, PBS Only

Catalog Number: 86226-3-PBS



## Basic Information

|                                     |                                                             |                                                |
|-------------------------------------|-------------------------------------------------------------|------------------------------------------------|
| Catalog Number:<br>86226-3-PBS      | GenBank Accession Number:<br>BC028050                       | Purification Method:<br>Protein A purification |
| Concentration:<br>1000 µg/ml        | GeneID (NCBI):<br>23373                                     | CloneNo.:<br>250870G10                         |
| Source:<br>Rabbit                   | UNIPROT ID:<br>Q6UUV9                                       |                                                |
| Isotype:<br>IgG                     | Full Name:<br>CREB regulated transcription<br>coactivator 1 |                                                |
| Immunogen Catalog Number:<br>AG0693 | Calculated MW:<br>75 kDa                                    |                                                |
|                                     | Observed MW:<br>68-80 kDa                                   |                                                |

## Applications

Tested Applications:  
WB, Indirect ELISA

Species Specificity:  
human

## Background Information

CRTC1, also named as MECT1, TORC1, and WAMP1, belongs to the TORC family. It is a transcriptional coactivator for CREB1 which activates transcription through both consensus and variant cAMP response element (CRE) sites. CRTC1 acts as a coactivator, in the SIK/TORC signaling pathway, being active when dephosphorylated and acts independently of CREB1 'Ser-133' phosphorylation. It enhances the interaction of CREB1 with TAF4. CRTC1 regulates the expression of specific CREB-activated genes such as the steroidogenic gene, StAR. It is a potent coactivator of PGC1alpha and inducer of mitochondrial biogenesis in muscle cells. CRTC1 is also a coactivator for TAX activation of the human T-cell leukemia virus type 1 (HTLV-1) long terminal repeats (LTR). In the hippocampus, CRTC1 involved in late-phase long-term potentiation (L-LTP) maintenance at the Schaffer collateral-CA1 synapses. It may be required for dendritic growth of developing cortical neurons. CRTC1 has some isoforms produced by alternative splicing with MW of 67 kDa, 68 kDa and 62 kDa. Sometimes it appears as a 75-80 kDa band probably due to phosphorylation (PMID: 17183642; 22770221)

## Storage

Storage:  
Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:  
PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 86226-3-RR (TORC1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 86226-3-PBS in a different storage buffer formulation.



Biolayer interferometry (BLI) kinetic assays of 86226-3-RR against Human TORC1/CRTC1 were performed. The affinity constant is below 1 pM.