

For Research Use Only

Acetyl-Histone H2B (Lys12) Recombinant monoclonal antibody, PBS Only

Catalog Number: 86731-1-PBS



Basic Information

Catalog Number:	86731-1-PBS	GenBank Accession Number:	BC005827	Purification Method:	Protein A purification
Source:	Rabbit	GeneID (NCBI):	8349	CloneNo.:	250005E8
Isotype:	IgG	UNIPROT ID:	Q16778		
		Full Name:	histone cluster 2, H2be		
		Calculated MW:	14 kDa		
		Observed MW:	15 kDa		

Applications

Tested Applications:
WB, IHC, FC (Intra), Dot blot, Indirect ELISA, ChIP-qPCR

Species Specificity:
human, mouse

Background Information

Histones are small, highly basic proteins that consist of a globular domain with unstructured N- and C-terminal tails protruding from the main structure. Histone H3 is one of the five main histones that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. In addition to their role in DNA compartmentalization, histones also play crucial roles in various biologic processes, including gene expression and regulation, DNA repair, chromatin condensation, cell cycle progression, chromosome segregation, and apoptosis. The ability of histones to regulate chromatin dynamics primarily originates from various posttranslational modifications carried out by histone-modifying enzymes.

Storage


Storage:
Store at -80°C.
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:
PBS only, pH7.3


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Dot blot analysis was used to confirm the specificity of 86731-1-RR Acetyl-Histone H2B (Lys12) antibody. peptides were spotted onto NC and probed with antibody at 1 µg/ml. The amount of peptide (µ g/mL) spotted is indicated next to each row. This data was developed using the same antibody clone with 86731-1-PBS in a different storage buffer formulation.




Chromatin was prepared from HeLa cells. Cells were fixed with formaldehyde for 10 minutes. The ChIP was performed with 15 µg of cross-linked chromatin, 5 µg of Acetyl-Histone H2B (Lys12) (86731-1-RR) or 5 µg of Normal Rabbit IgG (98136-1-RR), and 20 µl of Protein A Magarose Beads. The immunoprecipitated DNA was quantified by real-time PCR. This data was developed using the same antibody clone with 86731-1-PBS in a different storage buffer formulation.




Trichostatin A treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 86731-1-RR (Acetyl-Histone H2B (Lys12) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with GAPDH Monoclonal antibody (60004-1-Ig) as loading control. This data was developed using the same antibody clone with 86731-1-PBS in a different storage buffer formulation.




Trichostatin A treated HeLa cells were subjected to SDS PAGE followed by western blot with 86731-1-RR (Acetyl-Histone H2B (Lys12) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with GAPDH Monoclonal antibody (60004-1-Ig) as loading control. This data was developed using the same antibody clone with 86731-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 86731-1-RR (Acetyl-Histone H2B (Lys12) antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 86731-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 86731-1-RR (Acetyl-Histone H2B (Lys12) antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 86731-1-PBS in a different storage buffer formulation.



1x10⁶ Trichostatin A treated NIH/3T3 cells were intracellularly stained with 0.13 ug Acetyl-Histone H2B (Lys12) Recombinant monoclonal antibody (86731-1-RR, Clone:250005E8) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.13 ug Isotype Control (blue). Cells were fixed and permeabilized with Flow Cytometry Phosphorylated Protein Fixation/Permeabilization Kit (PF00026). This data was developed using the same