

For Research Use Only

# Di-Methyl-Histone H3 (Lys79) Recombinant monoclonal antibody, PBS Only

Catalog Number: 86734-1-PBS



## Basic Information

<b>Catalog Number:</b> 86734-1-PBS	<b>GenBank Accession Number:</b> BC066245	<b>Purification Method:</b> Protein A purification
<b>Source:</b> Rabbit	<b>GeneID (NCBI):</b> 8350	<b>CloneNo.:</b> 251570E11
<b>Isotype:</b> IgG	<b>UNIPROT ID:</b> P68431	
	<b>Full Name:</b> histone cluster 1, H3a	
	<b>Observed MW:</b> 15 kDa	

## Applications

**Tested Applications:**  
WB, Dot Blot, Indirect ELISA, ChIP-qPCR

**Species Specificity:**  
human, mouse

## Background Information

Histones are small, highly basic proteins that consist of a globular domain with unstructured N- and C-terminal tails protruding from the main structure. Histone H3 is one of the five main histones that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. In addition to their role in DNA compartmentalization, histones also play crucial roles in various biologic processes, including gene expression and regulation, DNA repair, chromatin condensation, cell cycle progression, chromosome segregation, and apoptosis. The ability of histones to regulate chromatin dynamics primarily originates from various posttranslational modifications carried out by histone-modifying enzymes.

## Storage

**Storage:**  
Store at -80°C.  
**The product is shipped with ice packs. Upon receipt, store it immediately at -80°C**

**Storage Buffer:**  
PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




Dot blot analysis was used to confirm the specificity of 86734-1-RR Di-Methyl-Histone H3 (Lys79) antibody. peptides were spotted onto NC and probed with antibody at 1  $\mu\text{g/mL}$ . The amount of peptide ( $\mu\text{g/mL}$ ) spotted is indicated next to each row. This data was developed using the same antibody clone with 86734-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 86734-1-RR (Di-Methyl-Histone H3 (Lys79) antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 86734-1-PBS in a different storage buffer formulation.



Chromatin was prepared from HeLa cells. Cells were fixed with formaldehyde for 10 minutes. The ChIP was performed with 15  $\mu\text{g}$  of cross-linked chromatin, 5  $\mu\text{g}$  of Di-Methyl-Histone H3 (Lys79) (86734-1-RR) or 5  $\mu\text{g}$  of Normal Rabbit IgG (88136-1-RR), and 20  $\mu\text{l}$  of Protein A Magarose Beads. The immunoprecipitated DNA was quantified by real-time PCR. This data was developed using the same antibody clone with 86734-1-PBS in a different storage buffer formulation.



mouse liver tissue were subjected to SDS PAGE followed by western blot with 86734-1-RR (Di-Methyl-Histone H3 (Lys79) antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 86734-1-PBS in a different storage buffer formulation.