

For Research Use Only

Phospho-CAD (Ser1859) Recombinant monoclonal antibody, PBS Only

Catalog Number: 87337-2-PBS



Basic Information

Catalog Number:

87337-2-PBS

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_004341

GeneID (NCBI):

790

UNIPROT ID:

P27708

Full Name:

carbamoyl-phosphate synthetase 2, aspartate transcarbamylase, and dihydroorotase

Calculated MW:

2225 aa, 243 kDa

Observed MW:

250 kDa

Purification Method:

Protein A purification

CloneNo.:

252640C5

Applications

Tested Applications:

WB, IF/ICC, Indirect ELISA

Species Specificity:

human

Background Information

CAD (carbamoyl-phosphate synthetase 2, aspartate transcarbamylase, and dihydroorotase) is a multifunctional enzyme required for the de novo synthesis of pyrimidine nucleotides. Phospho-CAD (Ser1859) refers to the phosphorylated form of the multi-enzyme CAD (carbamoyl-phosphate synthetase 2, aspartate transcarbamylase, and dihydroorotase) on serine 1859. This modification is catalyzed by ribosomal protein S6 kinase 1 (S6K1) downstream of mTORC1 signaling and serves as a metabolic switch that converts CAD from its basal state into a highly active oligomer, thereby accelerating the first three rate-limiting steps of de novo pyrimidine synthesis. (PMID:25422319, PMID: 32974014)

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Non-treated HEK-293 cells, phosphatase inhibitor treated HEK-293 cells and λ phosphatase treated HEK-293 cells were subjected to SDS PAGE followed by western blot with 87337-2-RR (Phospho-CAD (Ser1859) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Vinculin (86930-1-RR) antibody as a loading control. This data was developed using the same antibody clone with 87337-2-



Non-treated HeLa cells, phosphatase inhibitor treated HeLa cells and λ phosphatase treated HeLa cells were subjected to SDS PAGE followed by western blot with 87337-2-RR (Phospho-CAD (Ser1859) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Vinculin (86930-1-RR) antibody as a loading control. This data was developed using the same antibody clone with 87337-2-



Non-treated MCF-7 cells and Rapamycin treated MCF-7 cells were subjected to SDS PAGE followed by western blot with 87337-2-RR (Phospho-CAD (Ser1859) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Vinculin (86930-1-RR) antibody as a loading control. This data was developed using the same antibody clone with 87337-2-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Methanol) fixed Rapamycin treated MCF-7 cells using Phospho-CAD (Ser1859) antibody (87337-2-RR, Clone: 252640C5) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). This data was developed using the same antibody clone with 87337-2-PBS in a different storage buffer formulation.