

For Research Use Only

Mouse CDCP1/CD318 Recombinant monoclonal antibody, PBS Only

Catalog Number: 87514-4-PBS



Basic Information

Catalog Number: 87514-4-PBS	GenBank Accession Number: NM_133974.3	Purification Method: Protein A purification
Source: Rabbit	GeneID (NCBI): 109332	CloneNo.: 251667H4
Isotype: IgG	UNIPROT ID: Q5U462	
Immunogen Catalog Number: EG6199	Full Name: CUB domain containing protein 1	
	Calculated MW: 93 kDa	
	Observed MW: 130-140 kDa	

Applications

Tested Applications:
WB, Indirect ELISA

Species Specificity:
mouse, rat

Background Information

CD318 (also known as CDCP1, TRASK, SIMA135, or gp140) is a cell-surface glycoprotein with an apparent molecular mass of 140 kDa. It consists of three extracellular CUB domains, a transmembrane domain, and an intracellular domain. CD318 can be proteolytically cleaved between the two distal CUB domains by certain serine proteases, resulting in different ratios of the 140-kDa intact molecule and the 80-kDa cleaved product on various cells. Cleaved CD318 is phosphorylated and activated by Src kinase, then the activated CD318 forms a complex with activated β 1 integrin and activates FAK/PI3K/Akt motility signaling to promote early tumor dissemination. Under normal conditions, CD318 is present on many epithelial cells, some hematopoietic cells (18), and mesenchymal stem cells. CD318 is also present on many tumor cells. Up-regulation of CD318 expression is associated with a poor prognosis for many cancer patients.


Storage

Storage:
Store at -80°C.
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C
Storage Buffer:
PBS only, pH7.3


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 87514-4-RR (CDCP1/CD318 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 87514-4-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 87514-4-RR (CDCP1/CD318 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 87514-4-PBS in a different storage buffer formulation.