

For Research Use Only

Pig IgD Recombinant monoclonal antibody

Catalog Number: 87737-1-RR



Basic Information

Catalog Number: 87737-1-RR	GenBank Accession Number: /	Purification Method: Protein A purification
Source: Rabbit	GeneID (NCBI): /	CloneNo.: 253406C11
Isotype: IgG	UNIPROT ID: K7ZRJ8	Recommended Dilutions: WB: 1:1000-1:6000
Immunogen Catalog Number: EG7410	Full Name: Calculated MW: 43 kDa Observed MW: 75 kDa and 25 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls:
Species Specificity: pig	WB : pig spleen tissue, pig serum sample

Background Information

Immunoglobulin D (IgD) is an antibody class primarily functioning as a membrane-bound receptor on the surface of naïve B cells. Comprising less than 1% of serum immunoglobulins, IgD exists mainly in a transmembrane form that co-expresses with IgM on mature B lymphocytes. It plays a crucial role in B cell activation, survival, and immune surveillance by sensing environmental antigens and regulating humoral immune responses. Recent studies reveal IgD's involvement in mucosal immunity and its ability to bind to basophils and mast cells, triggering antimicrobial defense. Structurally, IgD features an extended hinge region, making it susceptible to proteolytic cleavage. Despite being discovered in 1965, IgD's precise functions remain an active area of immunological research. The immunogen for this antibody is Eg7410, and the immunogen sequence targets the Fc region of porcine IgD.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 87737-1-RR (Pig IgD antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLI) kinetic assays of 87737-1-RR against Pig IgD were performed. The affinity constant is 0.22 nM.