

For Research Use Only

ACAA1 Recombinant monoclonal antibody

Catalog Number:87969-1-RR



Basic Information

Catalog Number: 87969-1-RR	GenBank Accession Number: BC011977	Purification Method: Protein A purification
Source: Rabbit	GeneID (NCBI): 30	CloneNo.: 260194F9
Isotype: IgG	UNIPROT ID: P09110	Recommended Dilutions: WB: 1:5000-1:50000
Immunogen Catalog Number: AG2972	Full Name: acetyl-Coenzyme A acyltransferase 1	
	Calculated MW: 424 aa, 44 kDa	
	Observed MW: 41 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : A549 cells, SK-BR-3 cells, U-251 cells, HuH-7 cells, mouse liver tissue, rat liver tissue
Species Specificity: human, mouse, rat	

Background Information

ACAA1 (acetyl-CoA acyltransferase 1) is an enzyme involved in lipid β -oxidation and provides substrates to the tricarboxylic acid (TCA) cycle, a critical step in cellular metabolism (PMID:33194642). ACAA1 is also a biomarker in type 2 diabetes (T2D), predicting the pre-diabetic metabolic signature in mouse models (PMID:25784953). Moreover, ACAA1 is highly expressed in TNBC, serving as a potential therapeutic target in ACAA1-high tumors and a predictive biomarker of resistance to CDK4/6 inhibitors for RB1-proficient patients (PMID:37129951).

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 87969-1-RR (ACAA1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 87969-1-RR (ACAA1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 87969-1-RR (ACAA1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.