

For Research Use Only

# Frizzled 4 Recombinant monoclonal antibody

Catalog Number:87975-1-RR



## Basic Information

Catalog Number:

87975-1-RR

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

EG5015

GenBank Accession Number:

BC114527

GeneID (NCBI):

8322

UNIPROT ID:

Q9ULV1

Full Name:

frizzled homolog 4 (Drosophila)

Observed MW:

60 kDa

Purification Method:

Protein A purification

CloneNo.:

260031F12

Recommended Dilutions:

WB: 1:5000-1:50000

## Applications

Tested Applications:

WB, ELISA

Species Specificity:

human, mouse, rat

Positive Controls:

WB : mouse eye tissue, rat eye tissue, mouse heart tissue, rat heart tissue, human skeletal muscle tissue, human heart tissue, human adipose tissue

## Background Information

Frizzled 4, officially known as frizzled class receptor 4 and designated FZD4, is a protein encoded by the FZD4 gene in humans. It is a member of the frizzled (Fz) gene family, which encodes seven-transmembrane domain proteins that primarily function as receptors for Wnt signaling proteins. Frizzled 4 is embedded in the outer membrane of many cell types, where it plays a critical role in transmitting chemical signals from outside the cell to the nucleus. Frizzled 4 has an almost ubiquitous expression pattern and is found in a wide range of human tissues. Mutations in the FZD4 gene are linked to familial exudative vitreoretinopathy (EVR1), an eye disorder that affects the retina and can lead to progressive vision loss.

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)

W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 87975-1-RR (FZD4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.