

For Research Use Only

SLC6A8 Recombinant monoclonal antibody, PBS Only

Catalog Number: 88049-1-PBS



Basic Information

Catalog Number:

88049-1-PBS

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG14500

GenBank Accession Number:

BC012355

GeneID (NCBI):

6535

UNIPROT ID:

P48029

Full Name:

solute carrier family 6
(neurotransmitter transporter,
creatine), member 8

Calculated MW:

635 aa, 71 kDa

Observed MW:

65-75 kDa

Purification Method:

Protein A purification

CloneNo.:

260302B8

Applications

Tested Applications:

WB, Indirect ELISA

Species Specificity:

human, mouse, rat

Background Information

SLC6A8, also known as the sodium- and chloride-dependent creatine transporter 1 (CT1), plays a critical role in transporting creatine, a crucial molecule for energy metabolism, into cells. SLC6A8 belongs to the solute carrier family 6 (SLC6), responsible for transporting diverse molecules across cell membranes. SLC6A8 expression is highest in muscle, kidney, and other tissues with high energy demands. Mutations in SLC6A8 cause creatine transporter deficiency, an X-linked mental retardation disorder (PMID: 17465020). The 75 kDa band is a glycosylated modified form.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



rat kidney tissue were subjected to SDS PAGE followed by western blot with 88049-1-RR (SLC6A8 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 88049-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 88049-1-RR (SLC6A8 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 88049-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 88049-1-RR (SLC6A8 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 88049-1-PBS in a different storage buffer formulation.