Anti-Human CD200 Rabbit Recombinant Antibody, PBS Only

Antibodies | ELISA kits | Proteins WWW.ptglab.com


Catalog Number:98007-1-PBS

Catalog Number: 98007-1-PBS	GenBank Accession Number: BC022522	Purification Method: Protein A purification
Size: 1mg, 2000 µg/ml	GenelD (NCBI): 4345	CloneNo.: 230268E6
Source: Rabbit	UNIPROT ID: P41217	
Isotype: IgG	Full Name: CD200 molecule	
	Calculated MW: 269 aa, 30 kDa	
Tested Applications: FC		
Species Specificity: Human		
CD200, also known as OX2, is a type I transmembrane glycoprotein that belongs to the immunoglobulin superfamily (PMID: 3032785). It contains two extracellular immunoglobulin domains, a transmembrane and a cytoplasmic domain. CD200 is expressed on a variety of cell types, including thymocytes, B lymphocytes, a subset of T lymphocytes, follicular dendritic cells, neurons, and endothelial cells (PMID: 3032785; 10981966). CD200 binds to its receptor, CD200R, which is primarily expressed by myeloid and T cell lineage (PMID: 15187158). The CD200-CD200R interaction plays a role in immunosuppression and suppression of anti-tumor immune responses (PMID: 22020332).		
Storage: Store at -80°C. Stable for one Storage Buffer: PBS only	e year after shipment.	
	Size: 1mg, 2000 µg/ml Source: Rabbit Isotype: IgG Tested Applications: FC Species Specificity: Human CD200, also known as OX2, i superfamily (PMID: 3032785 cytoplasmic domain. CD200 of T lymphocytes, follicular of to its receptor, CD200R, whic CD200R interaction plays a r 22020332). Storage: Storage Buffer:	 Size: GenelD (NCBI): 1mg, 2000 µ g/ml 4345 Source: UNIPROT ID: Rabbit P41217 Isotype: Full Name: IgG CD200 molecule Calculated MW: 269 aa, 30 kDa Tested Applications: FC Species Specificity: Human CD200, also known as OX2, is a type I transmembrane glycoprotein that superfamily (PMID: 3032785). It contains two extracellular immunoglod cytoplasmic domain. CD200 is expressed on a variety of cell types, incl of T lymphocytes, follicular dendritic cells, neurons, and endothelial ce to its receptor, CD200R, which is primarily expressed by myeloid and T CD200R interaction plays a role in immunosuppression and suppression 2020332). Storage: Storage Buffer:

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CD200 (98007-1-PBS, Clone:230268E6) or 0.25 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) and PE-conjugated Goat Anti-Rabbit IgG. Cells were then co-stained with APC Anti-Human CD19 (HIB19) (APC-65110, Clone: HIB19). Cells were not fixed. Lymphocytes were gated.