

For Research Use Only

# Anti-Mouse IL-17F Rabbit Recombinant Antibody

Catalog Number: 98019-1-RR



## Basic Information

<b>Catalog Number:</b> 98019-1-RR	<b>GenBank Accession Number:</b> NM_145856.2	<b>Purification Method:</b> Protein A purification
<b>Concentration:</b> 100ug, 1000 ug/ml	<b>GeneID (NCBI):</b> 257630	<b>CloneNo.:</b> 230026G6
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> Q7TNI7-1	
<b>Isotype:</b> IgG	<b>Full Name:</b> interleukin 17F	
	<b>Calculated MW:</b> 17KD	

## Applications

**Tested Applications:**  
FC (Intra)

**Species Specificity:**  
mouse

## Background Information

The interleukin 17 (IL-17) family of cytokines contains 6 structurally related cytokines, IL-17A, IL-17B, IL-17C, IL-17D, IL-17E and IL-17F. IL-17 family plays crucial roles in host defense against microbial organisms and in the development of inflammatory diseases. IL-17A is a pro-inflammatory cytokine that also has the capacity to promote angiogenesis and osteoclastogenesis. IL-17F shares the highest homology with IL-17A and signals via a receptor composed by the IL-17RA and IL-17RC subunits. IL-17A and IL-17F can form IL-17A/A or IL-17F/F homodimers, IL-17A/F heterodimers are also formed. IL-17A and IL-17F, produced by the Th17 CD4(+) T cell lineage, have been linked to a variety of inflammatory and autoimmune conditions. IL-17F levels are elevated in sera and lesional psoriatic skin compared to non-lesional tissue. IL-17F also has been implicated in the development of neutrophilic airway inflammation.

## Storage

**Storage:**  
Store at 2-8°C. Stable for one year after shipment.

**Storage Buffer:**  
PBS with 0.09% sodium azide.

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1x10<sup>6</sup> untreated or PMA, Ionomycin and Brefeldin A treated C57BL/6 Th17-polarized splenocytes were intracellularly stained with 0.25 ug Anti-Mouse IL-17F Rabbit Recombinant Antibody (98019-1-RR, Clone:230026G6) and PE-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). Cells were co-stained with CoraLite® Plus 647 Anti-Mouse CD4. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLI) kinetic assays of 98019-1-RR against Mouse IL-17F were performed. The affinity constant is below 1 pM.