## For Research Use Only

## Anti-Human TIGIT Rabbit Recombinant Antibody

Catalog Number: 98023-2-RR



**Basic Information** 

Catalog Number: 98023-2-RR

Concentration: 100ug, 1000 ug/ml

Source: Rabbit Isotype:

IgG

GenBank Accession Number:

NM\_173799.4 GeneID (NCBI): 201633 UNIPROT ID:

Q495A1-1 Full Name:

T cell immunor eceptor with  $\mbox{Ig}$  and

ITIM domains

Calculated MW:
26 kDa

Purification Method: Protein A purfication

CloneNo.: 240423C7

**Applications** 

**Tested Applications:** 

FC

Species Specificity:

human

## **Background Information**

TIGIT (T-cell immunoreceptor with Ig and ITIM domains), also known as VSIG9 or VSTM3, is an immune receptor expressed on T cells, including Treg and memory subsets, as well as on NK cells (PMID: 19011627). It contains an immunoglobulin variable domain, a transmembrane domain and an immunoreceptor tyrosine-based inhibitory motif (ITIM). TIGIT binds to poliovirus receptor (PVR, also called CD155) with high affinity, and also to PVRL2 (CD112) with lower affinity. The interaction of TIGIT with PVR on dendritic cells increases the secretion of IL-10 and decreases the secretion of proinflammatory cytokine and suppresses T-cell activation by promoting the generation of mature immunoregulatory dendritic cells. TIGIT can inhibit NK cytotoxicity directly through its ITIM (PMID: 19815499).

Storage


Storage:

Store at 2-8°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide.


## **Selected Validation Data**



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human TIGIT Rabbit Recombinant Antibody (98023-2-RR, Clone: 240423C7) or Isotype Control, and PE-conjugated Goat Anti-Rabbit IgG. Cells were then stained with APC Anti-Human CD45RO. Cells were not fixed. Lymphocytes were gated.



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human TIGIT Rabbit Recombinant Antibody (98023-2-RR, Clone: 240423C7) (red) or Isotype Control (blue), and PE-conjugated Goat Anti-Rabbit IgG. Cells were not fixed. Lymphocytes were gated.



Biolayer interferometry (BLI) kinetic assays of 98023-2-RR against Human TIGIT were performed. The affinity constant is 5.45 nM.