For Research Use Only

Anti-Mouse IFN-gamma Rabbit Recombinant Antibody

Catalog Number:98045-1-RR





Basic Information	Catalog Number: 98045-1-RR	GenBank Accession Number: BC119063	Purification Method: Protein A purfication
	Size: 100ug, 1000 ug/ml	Genel D (NCBI): 15978	CloneNo.: 240262C9
	Source: Rabbit	UNIPROT ID: P01580	
	lsotype: IgG	Full Name: interferon gamma	
		Calculated MW: 17 kDa	
Applications	Tested Applications: FC (Intra)		
	Species Specificity: mouse		
Background Information	Interferon gamma (IFN γ), is a type II interferon that provides immunity against bacterial, viral and protozoan infections. In its active form, IFN γ is a glycosylated, non-covalently linked homodimer of 29-32 kDa subunits. It is produced by a number of immune cell types including natural killer cells, natural killer T cells, and effector lymphocyte T cells following antigenic and inflammatory triggers. The IFN γ dimer binds to its cognate receptor which has two subunits: IFN- γ R1 which is the ligand-binding chain (α chain) and IFN- γ R2, the signal-transducing chain (β chain). Binding to the receptor activates the JAK/STAT pathway which in turn activates IFN γ responsive genes. While IFN γ can inhibit viral replication, it also works as an immune-modulator and immune-stimulator by increasing surface expression of class I MHC proteins. (PMID: 19268625; 10688427)		
Storage	Storage: Store at 2-8°C. Stable for one year at Storage Buffer: PBS with 0.09% sodium azide.	fter shipment.	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1x10^{^6} untreated or PMA, Ionomycin and Brefeldin A treated C57BL/6 Th1-polarized splenocytes were intracellularly stained with 0.25 ug Anti-Mouse IFN-gamma Rabbit Recombinant Antibody (98045-1-RR, Clone:240262C9) and PE-conjugated Goat Anti-Rabbit IgG. Cells were then stained with CoraLite® Plus 647 Anti-Mouse CD4. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).

Biolayer interferometry (BLI) kinetic assays of 98045-1-RR against Mouse IFN-gamma were performed. The affinity constant is below 1 pM.